

Seiberg-Witten theory and matrix models

Albrecht Klemm¹ and Piotr Sułkowski^{1,2}

¹ *Physikalisches Institut der Universität Bonn and Bethe Center for Theoretical Physics,
Nussallee 12, 53115 Bonn, Germany*

² *Soltan Institute for Nuclear Studies, ul. Hoża 69, 00-681 Warsaw, Poland*

aklemm@th.physik.uni-bonn.de, Piotr.Sulkowski@fuw.edu.pl

Abstract

We derive a family of matrix models which encode solutions to the Seiberg-Witten theory in 4 and 5 dimensions. Partition functions of these matrix models are equal to the corresponding Nekrasov partition functions, and their spectral curves are the Seiberg-Witten curves of the corresponding theories. In consequence of the geometric engineering, the 5-dimensional case provides a novel matrix model formulation of the topological string theory on a wide class of non-compact toric Calabi-Yau manifolds. This approach also unifies and generalizes other matrix models, such as the Eguchi-Yang matrix model, matrix models for bundles over $\mathbb{P}^1$, and Chern-Simons matrix models for lens spaces, which arise as various limits of our general result.

Contents

1	Introduction	3
2	Seiberg-Witten theory	5
2.1	Partitions and Plancherel measure	6
2.2	The Nekrasov partition function	7
2.3	Seiberg-Witten curve	7
2.4	5-dimensional generalization	9
3	Matrix model for 4-dimensional Seiberg-Witten theory	10
3.1	The Nekrasov partition function as a matrix model	10
3.2	Seiberg-Witten curve as a spectral curve	14
4	Matrix model for 5-dimensional gauge theory	15
4.1	5-dimensional partition function as a matrix model	16
4.2	5-dimensional spectral curve	19
5	Relation to geometric engineering and topological string theory	24
5.1	Geometric engineering of 5-dimensional gauge theories	25
5.2	Mirror symmetry for toric non-compact Calabi-Yau	26
5.3	Geometric engineering of 5-dimensional $SU(n)$ gauge theories	27
5.4	4-dimensional Seiberg-Witten limit	30
5.5	Matrix model description of the $m \neq 0$ cases	31
6	Relations to other matrix models	32
6.1	Chern-Simons matrix model for lens spaces	33
6.2	Matrix models for line bundles over $\mathbb{P}^1$	33
6.3	Eguchi-Yang matrix model	34
7	Conclusions and discussion	35

1 Introduction

Finding the solution of $\mathcal{N} = 2$ supersymmetric gauge theories by Seiberg and Witten in terms of associated families of hyperelliptic Riemann surfaces, the Seiberg-Witten curves [1, 2], was a significant development in theoretical physics. The so-called Seiberg-Witten theory unifies many branches of physics and mathematics. From the physics perspective generalizations to theories with various gauge groups and matter contents were studied, as well as their relation to and the embedding in string theory. From the mathematical viewpoint the Seiberg-Witten solution gave important insights into the topology of four-manifolds. Also the solubility of $\mathcal{N} = 2$ theories turned out to be related to the underlying integrability, and the relations to various integrable systems were found. The literature on all these developments is immense, and the good starting point might be to consult the following reviews [3, 4, 5, 6, 7, 8], as well as references therein.

The holomorphic partition function $\mathcal{Z}$ is a central object in the theory. Its asymptotic expansion $\mathcal{Z} = \exp \sum_{g=0} \hbar^{2g-2} \mathcal{F}_g(\mathbf{a}, \Lambda)$ in $\hbar$ is a generating function for gauge theory instanton numbers, which appear as coefficients of $\hbar$, and inverse powers of the vevs a_l in the diagonal of the adjoint Higgs field. In a_l^{-1} the $\mathcal{F}_g(\mathbf{a}, \Lambda)$ have a finite radius of convergence as it has to be the case for physical terms in the effective action. In particular $\mathcal{F} = \mathcal{F}_0(\mathbf{a}, \Lambda)$ is the prepotential, which determines the exact low energy gauge theory effective action up to two derivatives. The $\mathcal{F}_g(\mathbf{a}, \Lambda)$ for $g > 0$ multiply gravitational couplings of the form $R_+^2 F_+^{2g-2}$, where R_+ and F_+ are the self-dual parts of the curvature and the graviphoton field strength respectively.

The fact that $\mathcal{F}_0(\mathbf{a}, \Lambda)$ can be calculated from periods of a family of hyperelliptic Riemann surface $\Sigma(\mathbf{a}, \Lambda)$ over a meromorphic one form differential dS was the main insight of [1, 2]. However this was argued using global consistency conditions of the low energy effective action and not from the microscopic action itself. The latter argument was provided by Nekrasov and Okounkov [9, 10]. In [9] Nekrasov developed a direct instanton calculus and used localization techniques to determine $\mathcal{Z}$ to all orders in $\hbar$ as sums over two-dimensional partitions labeling the instanton configurations. In the thermodynamic limit the limiting shape of the partitions approaches the Seiberg-Witten curve $\Sigma(\mathbf{a}, \Lambda)$ and in [10] it was proven that the periods of the latter over dS reproduce $\mathcal{F}$ to all orders in a_l .

The curve $\Sigma(\mathbf{a}, \Lambda)$ and the differential dS are the defining data of the $\mathcal{N} = 2$ supersymmetric theories in the sense that $\mathcal{Z}$ can be reconstructed from them. In particular one can view $\Sigma(\mathbf{a}, \Lambda)$ as the spectral curve of a putative matrix model and the A-cycle integrals of

dS as the fixed filling fractions and use the recursive solutions [11] of the loop equation to reconstruct the $\mathcal{F}_g$.

In this paper we find the explicit 1-matrix model for $SU(n)$ Seiberg-Witten theory, whose spectral curve is the Seiberg-Witten curve. The strategy we use is to represent partitions by $N \times N$ matrix integrals in a way recently proposed by Eynard [14], which explicitly reproduces the Nekrasov partition function in the large N limit. The matrix model which we find reads

$$Z^{4d} = \int_{Mat_{nN}} \mathcal{D}M e^{-\frac{1}{\hbar} \text{Tr } V^{4d}(M)}, \quad \mathcal{D}M = \prod_i dx_i \prod_{i < j} (x_i - x_j)^2, \quad (1)$$

where $M \in Mat_{nN}$ is $nN \times nN$ matrix and the measure $\mathcal{D}M$ involves the ordinary Vandermonde determinant. In the large N limit the potential is given by

$$V^{4d}(x) = tx + 2 \sum_{l=1}^n \left((x - a_l) \log(x - a_l) - (x - a_l) \right),$$

where a_l are vevs of the Higgs field, while t in the linear term encodes in particular the scale Λ .

More precisely the fact that it is possible to write the Nekrasov partition function as a matrix integral is a consequence of its close relation to the Plancherel measure on partitions. The Nekrasov sum for $SU(n)$ theory can be regarded as a generalization of the Plancherel measure to the case of n sets of partitions; equivalently it can be written as sum over one set of the so-called blended partitions. The properties of the Plancherel measure have been known for a long time; in particular it is known that in the thermodynamic limit it leads to the smooth limiting shape of large partitions known as the arc-sin law [15, 16]. This was used by [10] to show that analogous limiting shapes arise for the case of the Nekrasov partition function, and that they encode the geometry of Seiberg-Witten curves of the corresponding $\mathcal{N} = 2$ theories. Then, recently Eynard demonstrated how to rewrite the ordinary Plancherel measure, as well as its q -deformation, as a matrix integral [14]. In this paper we use similar methods to derive the above matrix models for $SU(n)$ theory.

We also derive a matrix model for a 5-dimensional theory compactified on a circle. We find that it reads

$$Z^{5d} = \int_{Mat_{nN}} \mathcal{D}M e^{-\frac{1}{g_s} \text{Tr } V^{5d}(M)}, \quad \mathcal{D}M = \prod_i du_i \prod_{i < j} \left(2 \sinh \frac{u_i - u_j}{2} \right)^2, \quad (2)$$

which is related to a deformation of the 4-dimensional result. Now the measure is given

by the deformed Vandermonde determinant, while the potential reads

$$V^{5d}(u) = tu + \frac{n}{2}u^2 + 2 \sum_{l=1}^n \text{Li}_2(e^{u+a_l}). \quad (3)$$

As is well known, e.g. from the geometric engineering of gauge theories [18], the partition function for 5-dimensional theory is equal to the partition function of topological strings on appropriate geometry. Therefore our matrix models also provide a new formulation of topological string theory, in the large radius limit of geometries which admit a limit to $SU(n)$ theories. In particular, in the text we will provide matrix model expressions for various non-compact toric Calabi-Yau spaces.

A very important feature of the matrix models which we derive is the fact that in various limits they reduce to other well-known matrix models. In particular, the 4-dimensional model (which itself is a limiting case of the 5-dimensional model) is a direct generalization of the Eguchi-Yang matrix model [19, 20] and reduces to it for $n = 1$. The 5-dimensional model, also for $n = 1$, leads immediately to matrix models for line bundles over $\mathbb{P}^1$ [21]. On the other hand, in the so called orbifold limit which involves $t = 0$ and suppression of Li_2 terms, the 5-dimensional model becomes just the quadratic matrix model with the deformed potential, which is the Chern-Simons matrix model for lens spaces postulated in [22, 23] and analyzed in [24, 25]. These other seemingly unrelated matrix models turn out to be just various corners of a single matrix model landscape.

The paper is organized as follows. In section 2 we recall a necessary background on Seiberg-Witten theory and the Nekrasov partition function. In section 3 we derive a matrix model for 4-dimensional $\mathcal{N} = 2$ $SU(n)$ theory and show that its spectral curve coincides with the Seiberg-Witten curve of $SU(n)$ gauge theory. In section 4 a matrix model for 5-dimensional theory and its spectral curve are derived and analyzed. In section 5 we discuss its relation to topological string theory. In section 6 we recover other well-known matrix models as certain limits of our most general matrix model. Section 7 contains a discussion.

2 Seiberg-Witten theory

In this section we review the solution of the Seiberg-Witten theory in terms of two dimensional partitions, as well as the underlying family of curves, and set up a necessary notation. After introducing two-dimensional partitions and the Plancherel measure, we

discuss how its generalization leads to the Nekrasov-Okounkov partition function in 4 and then in 5 dimensions.

2.1 Partitions and Plancherel measure

A partition λ of an integer number $|\lambda|$ is a set of non-negative, non-increasing integers

$$\lambda = (\lambda_1, \lambda_2, \dots), \quad \lambda_1 \geq \lambda_2 \geq \dots \geq 0, \quad \sum_i \lambda_i = |\lambda|.$$

A partition can also be presented as a Young tableaux given by the set of aligned horizontal rows, each containing λ_i boxes. The number of rows of a partition is defined as the number of non-zero λ_i . When considering statistical ensembles of partitions one can choose various measures. A very important one is the Plancherel measure defined as

$$\begin{aligned} P(\lambda) &= \left(\frac{\dim \lambda}{|\lambda|!} \right)^2 = \frac{1}{\left(\prod_{i=1}^N (\lambda_i + N - i)! \right)^2} \prod_{i < j}^N (\lambda_i - \lambda_j - i + j)^2 = \\ &= \prod_{i < j}^{\infty} \left(\frac{\lambda_i - \lambda_j - i + j}{j - i} \right)^2 = \prod_{\square \in \lambda} \frac{1}{hook(\square)^2}, \end{aligned} \quad (4)$$

where $hook(\square)$ is the Hook-length of a given box in λ , N is the number of rows, and $\dim \lambda$ is the dimension of the representation of the symmetric group corresponding to the Young-Tableaux. In the thermodynamic limit with respect to the Plancherel measure the partitions approach a limiting shape given by the so-called arc-sin law [15, 16]. This shape is described by the arcsin function, hence its name. An important fact is that, after the rescaling leading to the smooth limiting shape, the limiting partition has a finite length. This phenomenon is called the arctic circle property.

One can also consider a q -deformation of the Plancherel measure, denoted P_q , which arises by replacing the integers in (4) by its q -deformation

$$[h] = q^{-h/2} - q^{h/2}, \quad [h]! = \prod_{i=1}^h (q^{-i/2} - q^{i/2}),$$

so that

$$P_q(\lambda) = \prod_{i < j} \left(\frac{[\lambda_i - \lambda_j - i + j]}{[j - i]} \right)^2. \quad (5)$$

2.2 The Nekrasov partition function

The partition function for $\mathcal{N} = 2$, $SU(n)$ theory was derived by Nekrasov in [9] and analyzed further in great detail in [10]. It is a generalization of the Plancherel measure to the case of n sets of partitions. It therefore can be written down as a sum over a set of n partitions $\lambda_i^{(l)}$, with $l = 1, \dots, n$ labeling various partitions and index $i \geq 0$ denoting i 'th row of a given partition. This partition function depends on the scalar vevs a_l via $a_{lk} = a_l - a_k$ and it reads [9, 10, 26, 27]

$$Z^{SU(n)} = \sum_{\vec{\lambda}=(\lambda^{(1)}, \dots, \lambda^{(n)})} \Lambda^{2n|\vec{\lambda}|} Z_{\vec{\lambda}}, \quad (6)$$

$$Z_{\vec{\lambda}} = \prod_{(l,i) \neq (k,j)} \frac{a_{lk} + \hbar(\lambda_i^{(l)} - \lambda_j^{(k)} + j - i)}{a_{lk} + \hbar(j - i)},$$

where $|\vec{\lambda}| = \sum_{l=1}^n |\lambda^{(l)}|$. The sums in the above expression are performed over all possible partitions. To convert this partition function into a matrix model expression, below we consider the sums over partitions with at most N non-zero rows. In this case Nekrasov partition function is recovered in large N limit, where $N \rightarrow \infty$ and 't Hooft coupling $t = \hbar N$ is fixed. More generally, we could introduce an independent number of rows N_l for each partition $\lambda^{(l)}$. However it would not affect the answer, because the final result is independent of N .

2.3 Seiberg-Witten curve

The solution of the Seiberg-Witten theory is encoded in the Seiberg-Witten curve. For $SU(n)$ gauge theory this is a hyperelliptic curve of genus $n - 1$, which can be defined by the equation

$$\Lambda^n \left(w + \frac{1}{w} \right) = P_n(x), \quad P_n(x) = x^n + u_1 x^{n-1} + \dots + u_n, \quad (7)$$

for $w, x \in \mathbb{C}$, which is related to the familiar hyperelliptic form $y^2 = P_n^2 - 4\Lambda^{2n}$ by the identification $w = \frac{1}{2\Lambda^n}(y + P_n)$. The A- and B-periods of the differential

$$dS = \frac{1}{2\pi i} x \frac{dw}{w}$$

of this curve correspond respectively to the vevs a_l and derivatives of the prepotential $\frac{1}{2\pi i} \frac{\partial \mathcal{F}_0}{\partial a_l}$.

Let us recall now how this curve emerges as the stationary point of functionals [10], which arise from the ADHM construction of the instanton moduli space [9]. The k instanton partition function can be represented by contour integrals w.r.t. certain ADHM eigenvalues ϕ_I

$$Z_k = \sum_{|\vec{\lambda}|=k} Z_{\vec{\lambda}} = \oint \prod_{I=1}^k \left[\frac{\epsilon_1 + \epsilon_2}{2\pi\epsilon_1\epsilon_2} \frac{d\phi_I}{P(\phi_I)P(\phi_I + \epsilon_1 + \epsilon_2)} \right] \prod_{I \neq J} \frac{\phi_{IJ}(\phi_{IJ} + \epsilon_1 + \epsilon_2)}{(\phi_{IJ} + \epsilon_1)(\phi_{IJ} + \epsilon_2)}.$$

Here we wrote a more general expression with ϵ_1, ϵ_2 equivariant parameters, which lead to the Seiberg-Witten solution under the identification $\hbar = \epsilon_1 = -\epsilon_2$ (which we also assume in this paper). Moreover $\phi_{IJ} = \phi_I - \phi_J$ and

$$P(x) = \prod_{l=1}^n (x - a_l). \quad (8)$$

One can now introduce the density of eigenvalues

$$\rho(x) = \epsilon_1\epsilon_2 \sum_{I=1}^k \delta(x - \phi_I),$$

in terms of which the measure is approximated for small ϵ_1, ϵ_2 by the saddle point method as

$$\Lambda^{2kn} Z_k \sim \exp \left(\frac{1}{\epsilon_1\epsilon_2} \mathbf{E}_{\Lambda}[\rho] \right)$$

with ρ the saddle point of the functional

$$\mathbf{E}[\rho] = - \int^{p.v.} dx dy \frac{\rho(x)\rho(y)}{(x-y)^2} - 2 \int dx \rho(x) \log P(x). \quad (9)$$

Moreover, under the identification

$$\rho(x) = f(x) - \sum_{l=1}^n |x - a_l|,$$

the above functional is equivalent to the functional

$$\begin{aligned} \mathcal{E}[f] &= \frac{1}{4} \int_{y < x}^{p.v.} dx dy f''(x) f''(y) (x-y)^2 \left(\log(x-y) - \frac{3}{2} \right) = \\ &= -\frac{1}{2} \int_{x < y}^{p.v.} dx dy (N + f'(x))(N - f'(y)) \log(y-x), \end{aligned} \quad (10)$$

whose stationary point corresponds to the partition which arises as a limit shape f_* of Young diagrams in (6). This limit shape partition for $SU(n)$ theory is given by the function $f(x)$ of the form

$$f(x) = \sum_{l=1}^n f_l(x - a_l).$$

In fact, the stationary point f_* of the functional $\mathcal{E}[f]$ is related to the function $\varphi(x)$

$$f'_*(x) = \text{Re } \varphi(x), \quad (11)$$

and this is $\varphi(x)$ which explicitly encodes the Seiberg-Witten curve. More precisely, $\varphi(x) = \Phi(x + i0)$, where Φ is a certain conformal mapping described in detail in [10], such that the so-called bands of Φ determine the cuts of the Seiberg-Witten curve. The above construction was extended in [28], where the limit shape equations were cast in the form of the eigenvalue distribution for all classical gauge groups.

2.4 5-dimensional generalization

So far we considered the 4-dimensional theory. It can be regarded as a limit of a 5-dimensional gauge theory compactified on a circle of circumference β . For finite β the quantities in 5-dimensional theory, such as the partition function, correlation functions, or the spectral curve, are trigonometric analogues of the corresponding 4-dimensional quantities. In particular the partition function reads [9, 10, 26, 27]

$$Z_{5d}^{SU(n)} = \sum_{\vec{\lambda}=(\lambda^{(1)}, \dots, \lambda^{(n)})} (\beta\Lambda)^{2n|\vec{\lambda}|} Z_{\beta, \vec{\lambda}}, \quad (12)$$

$$Z_{\beta, \vec{\lambda}} = \prod_{(l,i) \neq (k,j)} \frac{\sinh \frac{\beta\hbar}{2}(p_{lk} + \lambda_i^{(l)} - \lambda_j^{(k)} + j - i)}{\sinh \frac{\beta\hbar}{2}(p_{lk} + j - i)}, \quad (13)$$

where $|\vec{\lambda}| = \sum_{l=1}^n |\lambda^{(l)}|$. Here and in what follows we often use the (quantized) values $p_l = a_l/\hbar$, and $p_{lk} = p_l - p_k$.

This theory is again characterized by a Riemann surface and a meromorphic differential. The corresponding limit shape can be found from the variational problem for the following functional [10]

$$\mathcal{E}_\beta[f] = \int_{x < y}^{p.v.} dx dy (N + f'(x))(N - f'(y)) \log \left(\frac{2}{\beta} \sinh \frac{\beta|y - x|}{2} \right), \quad (14)$$

which is a direct generalization of (10). In section 4.2 we will show that the corresponding curve can be found in a complementary way in the matrix model formalism, using the Migdal-Muskhelishvili formula for the resolvent. The relation of 5-dimensional gauge theory to topological string on non-compact Calabi-Yau spaces will also be discussed in section 5.

3 Matrix model for 4-dimensional Seiberg-Witten theory

In this section we derive matrix models for 4-dimensional Seiberg-Witten theory. Our strategy is to introduce an auxiliary dependence on a size N of two-dimensional partitions in the Nekrasov partition function, and replace the sum over these partitions by a sum over $N \times N$ matrices using the techniques presented in [14]. This leads to a matrix model with a very complicated potential, which includes such terms as logarithms of the Γ -function. However the result we are after is rederived in the $N \rightarrow \infty$ limit, which turns into the ordinary large N limit in the matrix model formalism. In this limit we get a matrix model with a smooth, (relatively) nice and (relatively) familiar potential. We also discuss how the Seiberg-Witten curve arises as a spectral curve of this matrix model.

3.1 The Nekrasov partition function as a matrix model

Our aim is to rewrite the partition function of the Seiberg-Witten theory (6) as a matrix model. For simplicity, we consider first $SU(2)$ case with $n = 2$. We denote $\lambda = \lambda^{(1)}$, $\mu = \lambda^{(2)}$, and introduce

$$h_i = \lambda_i - i + N + p_1, \quad k_i = \mu_i - i + N + p_2, \quad (15)$$

where we can assume $p_1 \ll p_2$, so that

$$k_1 > k_2 > \dots > k_N \geq p_2 > h_1 > h_2 \dots > h_N \geq p_1. \quad (16)$$

In terms of the variables h_i, k_j in the above range the $SU(2)$ partition function can be rewritten as

$$Z^{SU(2)} \sim \sum_{(h_i), (k_j)} \Lambda^{2n \sum_i (h_i + k_i)} \left(\prod_{i < j} \frac{h_i - h_j}{j - i} \right)^2 \left(\prod_{i < j} \frac{k_i - k_j}{j - i} \right)^2 \left(\prod_{i, j} \frac{h_i - k_j}{p_{12} + j - i} \right)^2.$$

The products in the above expression can be reorganized so that only non-trivial h_i, k_i with $i = 1, \dots, N$ occur explicitly. Furthermore the form of (15) leads for $i > N$ to many cancellations. Therefore we get for the first product above

$$\begin{aligned} \prod_{i < j} \frac{h_i - h_j}{j - i} &= \left(\prod_{1 \leq i < j \leq N} (h_i - h_j) \right) \left(\prod_{i=1}^N \prod_{j=1}^{\infty} \frac{h_i + j - p_1}{j} \right) = \\ &= \left(\prod_{1 \leq i < j \leq N} (h_i - h_j) \right) \left(\prod_{i=1}^N \frac{1}{(h_i - p_1)!} \right) \end{aligned}$$

and the analogous expression for the second product. The third product involves differences $(h_i - k_j)$ where the index i can be equal, smaller or greater than j . Similar cancellations as above lead to

$$\begin{aligned}
\prod_{i=j=1}^{\infty} \frac{h_i - k_j}{p_{12}} &= \left(\prod_{i=1}^N \frac{h_i - k_i}{p_{12}} \right) \left(\prod_{i=N+1}^{\infty} \frac{p_{12}}{p_{12}} \right) = \frac{1}{p_{12}^N} \prod_{i=1}^N (h_i - k_i) \\
\prod_{i < j} \frac{h_i - k_j}{p_{12} + j - i} &= (p_{12}!)^N \left(\prod_{1 \leq i < j \leq N} (h_i - k_j) \right) \left(\prod_{i=1}^N \frac{1}{(h_i - p_2)!} \right) \\
\prod_{i > j} \frac{h_i - k_j}{p_{12} + j - i} &= (-p_{12} + 1)_{p_{12}} \left(\prod_{1 \leq j < i \leq N} (h_i - k_j) \right) \left(\prod_{j=1}^N \frac{1}{(k_j - p_1)!} \right)
\end{aligned} \tag{17}$$

respectively.

Collecting all above contributions we observe that each difference $(h_i - h_j)$, $(k_i - k_j)$ and $(h_i - k_j)$ appears twice. There are also terms with factorials involving differences with p_1 and p_2 , which appear for both h_i and k_j on the same footing. We can therefore introduce one set of variables

$$l_{i=1, \dots, 2N} = (k_1, \dots, k_N, h_1, \dots, h_N). \tag{18}$$

In terms of l_i the partition function can be written in a compact form as

$$\begin{aligned}
Z^{SU(2)} &\sim N!^2 \sum_{l_1 > l_2 > \dots > l_N \geq p_2 > l_{N+1} > \dots > l_{2N} \geq p_1} \prod_{1 \leq i < j \leq 2N} (l_i - l_j)^2 \prod_{i=1}^{2N} \frac{\Lambda^{2nl_i}}{(l_i - p_1)!^2 (l_i - p_2)!^2} = \\
&= \sum_{l_1, \dots, l_N \geq p_2 > l_{N+1}, \dots, l_{2N} \geq p_1} \prod_{1 \leq i < j \leq 2N} (l_i - l_j)^2 \prod_{i=1}^{2N} \frac{\Lambda^{2nl_i}}{(l_i - p_1)!^2 (l_i - p_2)!^2},
\end{aligned} \tag{19}$$

which is normalized with respect to the constant factors like p_{12}^2 and $N!^2$; they are irrelevant for the further discussion and drop out anyway in correlation functions, thus can be safely skipped. In case when negatives quantities arise under the factorials in the above product, we replace these factorials by Γ functions, which provides the relevant analytic continuation.

The expression (19) already has a flavour of the matrix model, with the first product being the Vandermonde determinant. To write it explicitly as a matrix integral, we repeat the trick from [14] and introduce the function

$$f(x) = -x\Gamma(-x)\Gamma(x)e^{-i\pi x} = \frac{\pi e^{-i\pi x}}{\sin(\pi x)},$$

which has simple poles at all integer values of the argument. Upon integration along the contour C encircling $[p_1, \infty[$ part of the real axis, this function can be used to pick up all integer values of $h_i \in [p_1, \infty[$. Similarly, one would have to introduce another contour C' encircling $[p_2, \infty[$ half-line to pick up relevant values of k_i . In fact, we can use the same contour C for both h_i and k_j , because additional $k_j \in]p_1, p_2[$ picked up by the latter integral contribute zero (for such k_j , $\Gamma(k_j - p_2 + 1)^{-2}$ has a double zero which cancels the simple pole of $f(k_j)$). This leads to the following expression for (19)

$$\begin{aligned} Z^{SU(2)} &= \oint_{C^{2N}} dx_1 \cdots dx_{2N} \prod_{1 \leq i < j \leq 2N} (x_i - x_j)^2 \prod_{i=1}^{2N} \frac{f(x_i) \Lambda^{2nx_i}}{\Gamma(x_i - p_1 + 1)^2 \Gamma(x_i - p_2 + 1)^2} = \\ &= \int_{Mat_{2N}(C)} \mathcal{D}M e^{-\frac{1}{\hbar} \text{Tr } V(M)}, \end{aligned} \quad (20)$$

where $M \in Mat_{2N}(C)$ is $2N \times 2N$ matrix with eigenvalues in a set C . Choosing real values of p_l we can in fact reduce the contour to the real axis in the large N limit. It is convenient to rescale $x \mapsto x/\hbar$, and in terms of this rescaled variable the potential reads

$$\begin{aligned} \frac{1}{\hbar} V(x) &= -(2n \log \Lambda)x - \log f\left(\frac{x}{\hbar}\right) + \sum_{l=1,2} \log \Gamma(x\hbar^{-1} - p_l + 1)^2 = \\ &= -(2n \log \Lambda)x - \log\left(\frac{x}{\hbar}\right) - \log \Gamma\left(-\frac{x}{\hbar}\right) - \log \Gamma\left(\frac{x}{\hbar}\right) + \frac{i\pi x}{\hbar} + \\ &\quad + 2 \sum_l \left(\log(x\hbar^{-1} - p_l) + \log \Gamma(x\hbar^{-1} - p_l) \right). \end{aligned} \quad (21)$$

This expression looks quite complicated. Nonetheless, we will see that in the large N limit it simplifies enormously.

Now it also becomes clear that in generalization to arbitrary $SU(n)$ theory we need to introduce l_i for $i = 1, \dots, nN$, which is a concatenation of strings of N eigenvalues corresponding to matrices corresponding to each partition $\lambda^{(l)}$. In the last sum of (21) we also need to take the range of l from 1 to n .

In what follows also the derivative of this potential plays an important role. It involves the logarithmic derivative of the Γ function, i.e. the digamma function

$$\psi(x) = \frac{d}{dx} \log \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)}. \quad (22)$$

This function shares a number of interesting properties, for example

$$\psi(x+1) = -\gamma - \sum_{n=1}^{\infty} \left(\frac{1}{x+n} - \frac{1}{n} \right) = \log x + \frac{1}{2x} - \sum_{i=1}^{\infty} \frac{B_{2i}}{2ix^{2i}},$$

where γ is Euler-Mascheroni constant and B_{2i} are Bernoulli numbers. From this we also deduce

$$\psi(x) = -\frac{1}{2x} + \log x - \sum_{i=1}^{\infty} \frac{B_{2i}}{2ix^{2i}}.$$

From these expressions we find

$$\begin{aligned} V'(x) &= -2n\hbar \log \Lambda + \sum_l \left(\frac{1}{x\hbar^{-1} - p_l} + 2 \log \left(\frac{x}{\hbar} - p_l \right) - \sum_{i=1}^{\infty} \frac{B_{2i}}{i(x\hbar^{-1} - p_l)^{2i}} \right) = \\ &= -2n(\log \hbar + \hbar \log \Lambda) + \sum_l \left(\frac{\hbar}{x - a_l} + 2 \log(x - a_l) - \sum_{i=1}^{\infty} \frac{\hbar^{2i} B_{2i}}{i(x - a_l)^{2i}} \right). \end{aligned} \quad (23)$$

It is important to stress that the potential (21), and so of course its derivative above, explicitly do not depend on the size of matrices N .¹ For finite N , similarly as discussed in [14], there could be at most exponentially small corrections in N , which vanish in the large N limit we consider in what follows.

The above expression can also be integrated and yields

$$V(x) = tx + 2 \sum_l \left((x - a_l) \log(x - a_l) - (x - a_l) \right) + \mathcal{O}(\hbar), \quad (24)$$

where coefficients of linear terms are encoded in the constant t . We stress that so far we considered a finite value of N corresponding to the number of rows in partitions which appear in the Nekrasov partition function (6). Therefore the original Nekrasov partition function is automatically obtained in the large N limit for the matrix model expression (20)

$$N \rightarrow \infty, \quad \hbar \rightarrow 0, \quad \hbar N = \text{const}. \quad (25)$$

In this limit all the terms of order $\mathcal{O}(\hbar)$ vanish, so we conclude that the $SU(n)$ Seiberg-Witten theory can be formulated in terms of the 1-matrix model with the potential

$$V^{4d}(x) = tx + 2 \sum_{l=1}^n \left((x - a_l) \log(x - a_l) - (x - a_l) \right), \quad (26)$$

where t in the linear term encodes in particular the scale Λ . This potential is shown in figure 1. For $SU(n)$ theory it clearly has n minima to which we can associate n cuts, giving rise to the Seiberg-Witten spectral curve of genus $(n - 1)$.

¹Some complication due to explicit N dependence in eq. (2.23) in [14] arouse due to higher t_k couplings, which we do not consider here.




Figure 1: Matrix model potential for $SU(3)$ Seiberg-Witten theory. Left: the potential corresponding to the exact expression (21) for finite N ; complicated spikes arise from $\log \Gamma$ terms. Right: the potential in the large N limit, given by (26). Three minima give rise to three cuts and genus 2 Seiberg-Witten curve.

3.2 Seiberg-Witten curve as a spectral curve

Let us recall some basic facts concerning matrix models. For the matrix integral

$$Z = \int \mathcal{D}M e^{-\frac{1}{\hbar} \text{Tr} V(M)}$$

over $N \times N$ matrices, the eigenvalues are subject to the effective potential

$$S_{eff}(\lambda) = 2 \sum_{i < j} \log |\lambda_i - \lambda_j| + \frac{1}{\hbar} \sum_{i=1}^N V(\lambda_i).$$

Replacing discrete eigenvalues by a continuous eigenvalue density $r(x)$

$$\frac{1}{N} \sum_{i=1}^N h(\lambda_i) \mapsto \int h(x) r(x) dx$$

leads to the effective functional

$$S_{eff}[r] = \int_{x \neq y} dx dy r(x) r(y) \log |x - y| - \int dx r(x) V(x), \quad (27)$$

and the spectral curve of the matrix model is encoded in the discontinuities of the resolvent which leads to the minimization of this functional.

Let us now introduce $R(x)$ such that

$$R'(x) = r(x), \quad (28)$$

and integrate (27) by parts to get

$$S_{eff}[R] = \int_{x \neq y} dx dy \frac{R(x)R(y)}{(x-y)^2} + \int dx R(x)V'(x). \quad (29)$$

We focus now on the matrix model for Seiberg-Witten theory (20), with a potential whose derivative is given in (23). Let us stress that the matrix model (20) still depends on both N (size of a matrix) and $\hbar$. Upon extracting a spectral curve corresponding to the Seiberg-Witten theory we should again take the limit (25) in which all terms in (23) proportional to $\hbar$ are neglected. The only terms which survive are

$$V^{4d}(x) = 2 \sum_l \log(x - a_l) = 2 \log P(x), \quad (30)$$

where $P(x) = \prod_{l=1}^n (x - a_l)$ coincides with (8). Plugging this into (27) we get

$$S_{eff}[R] = \int_{x \neq y} dx dy \frac{R(x)R(y)}{(x-y)^2} + 2 \int dx R(x) \log P(x). \quad (31)$$

We observe that this functional is identical to (9) if we identify $R(x) = \rho(x)$. From the analysis in [10] we know that the stationary point of (9) leads to the Seiberg-Witten curve encoded in the derivative of the limiting profile (11). In our case we also need to take the derivative (28) of $R(x)$ to get $r(x)$ corresponding to the eigenvalue density. We conclude that the stationary point of (27) – encoding the spectral curve of our matrix model – corresponds to the same Seiberg-Witten curve as the one derived in [10].

The above argument relies on the form of the functional (9) considered by Nekrasov-Okounkov. There is however a way to obtain the spectral curve directly within the matrix model formalism. In more generality this works also for the 5-dimensional matrix model which we introduce in the next section. One can therefore derive the spectral curve for the 5-dimensional theory according to the procedure presented in the subsection 4.2, and then recover the curve discussed above in the 4-dimensional limit.

4 Matrix model for 5-dimensional gauge theory

In this section we generalize the construction of matrix models for 4-dimensional gauge theories in section 3 to a canonical class of 5-dimensional gauge theories. This construction will be further slightly generalized in section 5.

4.1 5-dimensional partition function as a matrix model

We now wish to introduce a matrix model for the 5-dimensional $SU(n)$ theory with a partition function given in (12). Similarly as before, to start with we consider $SU(2)$ case with $n = 2$. We denote $\lambda = \lambda^{(1)}$ and $\mu = \lambda^{(2)}$, which are partitions with no more than N rows, and similarly as in (15) introduce

$$h_i = \lambda_i - i + N + p_1, \quad k_i = \mu_i - i + N + p_2. \quad (32)$$

Moreover we assume $p_1 \ll p_2$, so that

$$k_1 > k_2 > \dots > k_N \geq p_2 > h_1 > h_2 \dots > h_N \geq p_1. \quad (33)$$

We also introduce $q = e^{-g_s} = e^{-\beta\hbar}$ and the standard q -deformed notation

$$[h] = q^{-h/2} - q^{h/2}, \quad [h]! = \prod_{i=1}^h (q^{-i/2} - q^{i/2}).$$

The $SU(2)$ partition function now takes the form

$$Z_{5d}^{SU(2)} \sim \sum_{(h_i), (k_j)} (\beta\Lambda)^{2n \sum_i (h_i + k_i)} \left(\prod_{i < j} \frac{[h_j - h_i]}{[i - j]} \right)^2 \left(\prod_{i < j} \frac{[k_j - k_i]}{[i - j]} \right)^2 \left(\prod_{i, j} \frac{[k_j - h_i]}{[i - j - p_{12}]} \right)^2,$$

with $(h_i), (k_j)$ satisfying the condition (33). These terms can be rewritten analogously as in section 3.1. For example

$$\prod_{i < j} \prod_{i < j} \frac{[h_j - h_i]}{[i - j]} = \left(\prod_{1 \leq i < j \leq N} [h_j - h_i] \right) \left(\prod_{i=1}^N \frac{1}{[h_i - p_1]!} \right),$$

and similarly for other terms. Therefore, introducing one set of variables

$$l_{i=1, \dots, 2N} = (k_1, \dots, k_N, h_1, \dots, h_N), \quad (34)$$

we get

$$Z_{5d}^{SU(2)} = \sum_{(l_i)_{i=1, \dots, 2N}} \left(\prod_{1 \leq i < j \leq 2N} [l_i - l_j]^2 \right) \left(\prod_{i=1}^{2N} \frac{(\beta\Lambda)^{2nl_i}}{([l_i - p_1]! [l_i - p_2]!)^2} \right). \quad (35)$$

To write this expression as a matrix model integral we again take advantage of a few facts discussed in [14]. First of all, we introduce the following notation for the quantum dilogarithm

$$g(x) = \prod_{i=1}^{\infty} (1 - x^{-1} q^i).$$

It vanishes $g(q^h) = 0$ for h a positive integer, and at such points its derivative is

$$g'(q^h) = -\frac{g(1)^2 e^{i\pi h} q^{-h(h-1)/2}}{q^h(1-q^h)g(q^{-h})}.$$

Therefore the following function has simple poles with residue 1 for $x = q^h$ with $h \in \mathbb{N}$

$$f(x) = -\frac{g(1)^2 e^{-\frac{i\pi}{g_s} \log x} e^{\frac{(\log x)^2}{2g_s}}}{(1-x)\sqrt{x}g(x)g(x^{-1})}.$$

Let us also note that for $x = q^l$

$$\begin{aligned} [l]! &= q^{-l(l+1)/4} \frac{g(1)}{g(x^{-1})}, \\ \frac{1}{([l-p]!)^2} &= \frac{g(x^{-1}q^p)^2}{g(1)^2} \exp\left(\frac{1}{2\log q} \log(xq^{-p}) \log(xq^{-p+1})\right). \end{aligned}$$

In what follows we use the notation

$$x_i = q^{l_i} = e^{-g_s l_i} = e^{u_i}, \quad (36)$$

so that x_i is a cylindrical coordinate. Now we can write

$$\begin{aligned} Z_{5d}^{SU(2)} &\sim \sum_{(l_i)_{i=1,\dots,2N}} \prod_{1 \leq i < j \leq 2N} (q^{l_i} - q^{l_j})^2 \prod_{i=1}^{2N} \frac{(\beta\Lambda)^{2nl_i} q^{(1-2N)l_i}}{([l_i - p_1]!)^2 ([l_i - p_2]!)^2} = \\ &= \int dx_1 \cdots dx_{2N} \prod_{1 \leq i < j \leq 2N} (x_i - x_j)^2 \times \\ &\quad \times \prod_{i=1}^{2N} \frac{(\beta\Lambda)^{-\frac{2n}{g_s} \log x_i} f(x_i)}{x_i^{2N-1}} \prod_{l=1}^{n=2} \left[\frac{g(x_i^{-1}q^{p_l})^2}{g(1)^2} e^{\frac{\log(x_i q^{-p_l}) \log(x_i q^{-p_l+1})}{2\log q}} \right]. \end{aligned} \quad (37)$$

The deformed Vandermonde determinant can also be written as

$$e^{(1-N) \sum_i \log x_i} \prod_{i < j} (x_i - x_j)^2 = \prod_{i < j} \left(2 \sinh \frac{\log x_i - \log x_j}{2} \right)^2 \equiv \Delta_q^2. \quad (38)$$

With the deformed measure

$$\mathcal{D}M = \Delta_q^2 \prod_i du_i = \Delta_q^2 e^{-\sum_i \log x_i} \prod_i dx_i$$

we can rewrite the above partition function as a matrix model expression, which in a straightforward way generalizes to arbitrary $SU(n)$ gauge group

$$Z_{5d}^{SU(n)} = \int_{M_{nN}(C_q)} \mathcal{D}M e^{-\frac{1}{g_s} \text{Tr } V_{5d}(M)},$$

with the deformed integration contour C_q is a circle of radius $r \in]1, q^{-1}[$. The matrix model potential takes the form

$$\begin{aligned} \frac{1}{g_s} V_{5d}(x) = & \left(\frac{2n \log(\beta\Lambda)}{g_s} + (n-1)N - 1 \right) \log x - 2 \log g(1) + \log g(x) + \log g(x^{-1}) + \\ & + \frac{1}{2} \log x + \log(1-x) + \frac{i\pi}{g_s} \log x - \frac{1}{2g_s} (\log x)^2 - i\pi + \\ & + \sum_{l=1}^n \left[2 \log g(1) - 2 \log g(x^{-1} q^{p_l}) - \frac{(\log(xq^{-p_l}))^2}{2 \log q} - \frac{1}{2} \log(xq^{-p_l}) \right]. \end{aligned} \quad (39)$$

Similarly as in 4-dimensional case, this expression also simplifies in the large N limit. To turn the potential into a nicer form, we again differentiate it and then integrate. First we note [14] that

$$\log g(x) = -\frac{1}{g_s} \sum_{m=0}^{\infty} \text{Li}_{2-m}(x^{-1}) \frac{B_m g_s^m}{m!},$$

where B_m are Bernoulli numbers. The polylogarithm is defined as

$$\text{Li}_m(x) = \sum_{i=1}^{\infty} \frac{x^i}{i^m},$$

and it has the following properties

$$\begin{aligned} \text{Li}'_n(x) &= \frac{\text{Li}_{n-1}(x)}{x}, \\ \text{Li}_m(x) &= (-1)^{m+1} \text{Li}_m(x^{-1}), \quad \text{for } m < 0 \\ \text{Li}_0(x) &= \frac{x}{1-x}, \\ \text{Li}_1(x) &= -\log(1-x). \end{aligned} \quad (40)$$

The following useful relations follow

$$\begin{aligned} \frac{g'(x)}{g(x)} &= \frac{1}{g_s x} \sum_{m=0}^{\infty} \text{Li}_{1-m}(x^{-1}) \frac{B_m g_s^m}{m!}, \\ x \frac{d}{dx} (\log g(x) + \log g(x^{-1})) &= \frac{\log x}{g_s} - \frac{1}{2} \frac{x+1}{x-1} - \frac{i\pi}{g_s}. \end{aligned}$$

Using the above facts, after some algebra one therefore finds the derivative of the potential

$$\begin{aligned} V'_{5d}(x) &= \frac{2n \log(\beta\Lambda) + g_s((n-1)N - \frac{1}{2})}{x} + \\ &+ \sum_{l=1}^n \left[\frac{g_s}{x - q^{p_l}} - \frac{2}{x} \log((xq^{-p_l})^{-\frac{1}{2}} - (xq^{-p_l})^{\frac{1}{2}}) + \frac{2}{x} \sum_{m=1}^{\infty} \text{Li}_{1-2m}(xq^{-p_l}) \frac{B_{2m} g_s^{2m}}{(2m)!} \right]. \end{aligned} \quad (41)$$

Let us note that $\log((xq^{-p_l})^{-\frac{1}{2}} - (xq^{-p_l})^{\frac{1}{2}})$ is the q -deformation of the 4-dimensional expression $\log(x - a_l)$ in (23), while other terms (apart from the linear in $1/x$) are proportional to higher powers of g_s . The expression which we obtained can now be integrated. In particular $\int \frac{dx}{x} \log(x^{-1/2} - x^{1/2}) = -\frac{1}{4}(\log x)^2 - \text{Li}_2(x)$, so that

$$V_{5d} = t \log x + \frac{n}{2}(\log x)^2 + 2 \sum_l \text{Li}_2(xe^{a_l}) + \mathcal{O}(g_s), \quad (42)$$

where the coefficient t encodes in particular the scale $\beta\Lambda$ and terms linear in a_l . In the large N limit

$$N \rightarrow \infty, \quad g_s \rightarrow 0, \quad g_s N = \text{const},$$

we can again neglect all terms $\mathcal{O}(g_s)$. We also note that introduction of the single parameter t is justified, as it involves both $\log(\beta\Lambda)$ and $g_s N = \text{const}$ related to each other in large N dualities precisely in this way. Substituting now $x = e^u$, finally we get the matrix model

$$Z_{5d}^{SU(n)} = \int \mathcal{D}M e^{-\frac{1}{g_s} \text{Tr } V^{5d}(M)} = \int \prod_i du_i \prod_{i < j} \left(2 \sinh \frac{u_i - u_j}{2} \right)^2 e^{-\frac{1}{g_s} \sum_i V^{5d}(u_i)} \quad (43)$$

with the potential

$$V^{5d}(u) = tu + \frac{n}{2}u^2 + 2 \sum_{l=1}^n \text{Li}_2(e^{u+a_l}). \quad (44)$$

The coefficient n in the quadratic term can be interpreted as redefining the coupling in front of the potential as

$$g_s \mapsto \hat{g}_s = \frac{g_s}{n}$$

(and rescaling other terms in the potential by n). It is consistent with the same rescaling found in a related matrix model in [23], which we discuss in section 6. The potential V^{5d} is shown in figure 2.

4.2 5-dimensional spectral curve

Finding the spectral curve for 5-dimensional theory also requires the 't Hooft limit, in which all terms in the potential (41) proportional to positive powers of g_s vanish, so that

$$xV'_{5d}(x) \longrightarrow -2 \sum_{l=1}^n \log((xq^{-p_l})^{-\frac{1}{2}} - (xq^{-p_l})^{\frac{1}{2}}) = -2 \log P_q(y),$$

$$\text{for } P_q(y) = \prod_{l=1}^n [y - p_l],$$




Figure 2: Matrix model potential for 5-dimensional $SU(3)$ Seiberg-Witten theory. Left: the potential in the large N limit, given by (44). Three minima give rise to three cuts and the genus 2 Seiberg-Witten curve. Right: zooming around the rightmost minimum.

with $x = q^y$. This is indeed the trigonometric counterpart of the 4-dimensional case (30). Accordingly the 5-dimensional curve will be the trigonometric counterpart of the 4-dimensional Seiberg-Witten curve, in agreement with the relation between 4-dimensional and the 5-dimensional variational problems (10) and (14).

Apart from arguments which rely on the form of effective functionals (which were also employed in section 3.2), we can in fact derive the form of the 5-dimensional spectral curve using purely matrix model technology, i.e. determining the resolvent and the eigenvalue density in the large N limit. This method could also have been applied explicitly in section 3.2, however it is more convenient to consider the general 5-dimensional case and then get the 4-dimensional results in the gauge theory limit. For a general potential in the multi-cut case the resolvent $\omega(z)$ is given by the Migdal-Muskhelishvili formula [29]. It has the following form in terms of the periodic variable x

$$\omega(z) = \frac{1}{2\pi i} \oint_C dx \frac{V'(x)}{x(z-x)} \sqrt{\prod_{k=1}^{2n} \frac{z-x_k}{x-x_k}}, \quad (45)$$

where $C = C_1 \cup \dots \cup C_n$ is a union of n cuts supported on intervals $C_k = [x_{2k-1}, x_{2k}]$. The $2n$ endpoints of the cuts x_k must be suitably chosen. A set of $(n+1)$ conditions for these endpoints stems from the limiting behaviour of the resolvent at infinity $\omega(z) \sim 1/z$. Another $(n-1)$ conditions can be chosen in a way which is appropriate for a given physical problem. In our case it is natural to fix $(n-1)$ filling fractions, which are determined by

the A-periods around the cuts.

To compute the resolvent (45) one can follow the solution presented in [21], where the case corresponding to $n = 1$ (which we discuss also in section 6.2) was analyzed in detail. This solution relies on the chiral ansatz and the arctic circle property of two-dimensional partitions which we discussed above. We assume that the eigenvalue density $\rho(z)$ is non-trivial only in certain intervals, independently for each set of partitions summed over in the Nekrasov partition function. The parameters a_i in the potential (44) can be absorbed into the positions of the endpoints of these integrals, so that the resolvent becomes

$$\omega(z) = \frac{1}{2\pi i} \oint_C dx \frac{n \log x + t - 2 \sum_{k=1}^n \log(1 - x e^{y_{2k}})}{x(z - x)} \sqrt{\prod_{k=1}^{2n} \frac{z - e^{-y_{2k-1}}}{x - e^{-y_{2k}}}}. \quad (46)$$

To perform this integral one can modify the integration contour, so that it encircles the branch cuts of all logarithms and the pole at $x = z$.




Figure 3: Integration contour used to compute the resolvent for 5-dimensional $SU(2)$ Seiberg-Witten theory.

As an example let us consider how to derive the spectral curve for the $SU(2)$ theory from our matrix model. We denote the endpoints of the two cuts as $[y_1, y_2]$ and $[y_3, y_4]$. The integration contour in this case is shown in figure 3. The resolvent becomes a sum of the residue at $x = z$ and the contour integral along the (infinitesimal) circle around $x = 0$ (a part of the contour around the branch cut of $\log x$), as well as contributions from the integrals along parts of real axis from branch cuts of all logarithms. The latter are given by integrals of the form

$$\int \frac{dv}{v(z - v) \sqrt{\prod_{i=1}^4 (v - e^{-y_i})}}. \quad (47)$$

They can be expressed in terms of elliptic functions of the first and the third kind, as explained in the appendix. In what follows we also use the notation introduced in the

appendix. After some simplifications, for $SU(2)$ theory the resolvent (46) turns out to involve three integrals of the form (47) evaluated at the points $v = 0, \infty, e^{-y_2}$ (the integral evaluated at e^{-y_4} vanish). For these points the arguments of the elliptic functions F and Π are respectively

$$y(0) = \frac{e^{-y_4}(e^{-y_3} - e^{-y_2})}{e^{-y_3}(e^{-y_4} - e^{-y_2})}, \quad y(\infty) = \frac{e^{-y_3} - e^{-y_2}}{e^{-y_4} - e^{-y_2}}, \quad y(e^{-y_2}) = 1,$$

as well as δ and k^2 given in (59) with a, b, c, d identified with e^{-y_i} . Using the addition formula (61) twice we can replace three elliptic functions $F(y(v), k)$ by a single one with the argument y_{0,∞,y_2} determined by using (64) twice. Similarly, using (62) twice we get a single function $\Pi(y_{0,\infty,y_2}, \delta, k)$, but then in addition two terms ($\text{artanh } \Theta_{0,\infty}$) and ($\text{artanh } \Theta_{0,\infty,y_2}$), where $\Theta_{0,\infty}$ and Θ_{0,∞,y_2} are determined from the formula (63) used twice. Altogether, the resolvent becomes

$$\begin{aligned} \omega(z) = & \frac{t + 2 \log z - 2 \log(1 - ze^{y_2}) - 2 \log(1 - ze^{y_4})}{z} + \frac{\sqrt{\prod_i (z - e^{-y_i})}}{z} t e^{\frac{y_1 + y_2 + y_3 + y_4}{2}} + \\ & + \frac{4 \sqrt{\prod_i (z - e^{-y_i})}}{z \sqrt{(e^{-y_4} - e^{-y_1})(e^{-y_3} - e^{-y_2})}} \left[\left(\frac{1}{e^{-y_3} - z} - \frac{1}{e^{-y_3}} \right) F(y_{0,\infty,y_2}) + \right. \\ & + \frac{e^{-y_4} - e^{-y_3}}{e^{-y_3} e^{-y_4}} \left(\Pi(y(0), \delta(0), k) + \Pi(y(\infty), \delta(0), k) - \Pi(y(e^{-y_2}), \delta(0), k) \right) + \\ & \left. + \frac{e^{-y_4} - e^{-y_3}}{(z - e^{-y_3})(z - e^{-y_4})} \left(\Pi(y_{0,\infty,y_2}, \delta(z), k) + \frac{\text{artanh } \Theta_{0,\infty} + \text{artanh } \Theta_{0,\infty,y_2}}{\sqrt{\delta(z)^{-1}(\delta(z) - 1)(\delta(z) - k^2)}} \right) \right]. \end{aligned}$$

We discuss now the behavior of this resolvent at infinity $z \rightarrow \infty$, where it should decay like $\sim 1/z$. Firstly, we note that the last term in the first line, the last term $\sim e^{y_3} F(y_{0,\infty,y_2})$ in the second line, and the terms in the third line are of order $\sim z$. Therefore a sum of coefficients of these terms must be zero, which gives one equation for the endpoints of the cuts. Due to simple the same dependence on z , these terms in fact cancel altogether and drop out from the expression. Then, the only term which is constant at infinity is the first term $\sim F(y_{0,\infty,y_2})/(e^{-y_3} - z)$ in the second line. It vanishes if and only if the elliptic function $F(y_{0,\infty,y_2})$ vanishes, which is possible if and only if $\sin \varphi \equiv y_{0,\infty,y_2} = 0$. On one hand, this provides another condition on the endpoints of the cuts, due to (64). On the other hand, then the terms in the fourth line $\Pi(y_{0,\infty,y_2} = 0, \delta(z), k)$, as well as ($\text{artanh } \Theta_{0,\infty,y_2}$) (whose argument is proportional to $\sin \varphi \equiv y_{0,\infty,y_2}$ via (63)) also vanish. Therefore the only term which could contribute to the eigenvalue density (or equivalently the spectral curve) is ($\text{artanh } \Theta_{0,\infty}$) in the fourth line. Quite non-trivial cancellations lead to its overall prefactor being just $2/z$ which in particular does not depend on any e^{-y_i} .

Moreover, this term indeed has the square root branching as hoped for, which arises via the dependence on z through the argument of $\Theta_{0,\infty}$

$$\sqrt{\delta(z)(1-\delta(z))(\delta(z)-k^2)} \sim \frac{\sqrt{\prod_i(z-e^{-y_i})}}{(e^{-y_4}-z)^2}.$$

Indeed some algebra reveals that $\Theta_{0,\infty} = \sqrt{\prod_i(z-e^{-y_i})}/P(z)$, with $P(z)$ a quadratic polynomial in z .

Identifying now the density with a new variable v/z we get

$$\frac{v}{z} \equiv \rho(z) = \frac{1}{2\pi i} (\omega(z-i\epsilon) - \omega(z+i\epsilon)) \sim \frac{2}{z} \operatorname{artanh} \Theta_{0,\infty} = \frac{1}{z} \log \frac{1+\Theta_{0,\infty}}{1-\Theta_{0,\infty}}.$$

Exponentiating both sides we get an equation in two $\mathbb{C}^*$ coordinates, z and $x \equiv e^v$. This defines a Riemann surface of genus one written in terms of two $\mathbb{C}^*$ coordinates x and z , as expected for 5-dimensional Seiberg-Witten theory. The latter one can be brought to hyperelliptic form after the redefinition $\frac{x-1}{x+1} \mapsto \tilde{x}$ and a rescaling of the new variable $\tilde{x}$ by $P(z)$, which to the equation of the curve

$$\tilde{x}^2 = \prod_{i=1}^4 (z - e^{-y_i}). \quad (48)$$

Three of the y_i 's are specified by the boundary conditions of the resolvent, and they depend also on the modulus t . To get the full solution, one more condition can be specified by imposing the A-period filling fraction. On the other hand, this A-period can be found from the knowledge of the above curve.

Let us stress at this point, that the appearance of integrals (47) of this form (in particular leading to elliptic functions for $SU(2)$ theory) is a consequence of the logarithmic branch cut structure of the derivative of the potential $V(x)$ which appears in (46). Derivatives of both the quadratic and dilogarithm terms which appear in the potential are logarithms with branch cuts that we just need. This is therefore non-accidental that no other (special) functions arise in the potential.

The above derivation could of course be generalized to arbitrary $SU(n)$ theory, though technically the solution would be even more complicated. In this general case the well-known matrix model relations [30] for the filling fractions a_l and the genus zero free energy F_0 , written in terms of the rescaled variables

$$a_l = \frac{1}{2\pi i} \oint_{A_l} \log x \frac{dz}{z}, \quad \frac{\partial F_0}{\partial a_l} = \frac{1}{2} \oint_{B_l} \log x \frac{dz}{z},$$

acquire a standard interpretation in Seiberg-Witten theory: they relate vevs of the Higgs field a_i with the prepotential F_0 . This completes then our solution of the Seiberg-Witten theory from the matrix model perspective.

We also note that the matrix models which we consider are unstable due to the unbounded potentials, as seen in figures 1 and 2. However this instability is not worse than the instability related to the condition of fixed filling fractions, which results in unequal level of a potential in each cut. Due to the condition of fixed filling fractions the eigenvalues are prevented from falling into the infinite instability well in the same way as they are prevented from tunnelling between cuts.

5 Relation to geometric engineering and topological string theory

The purpose of this section is to give an unified geometric description of the 4- and 5-dimensional supersymmetric gauge theories by geometric engineering of type II string theory and M-theory on non-compact Calabi-Yau spaces. This construction relates the matrix models discussed in this paper and the one in [22, 23, 24]. For 4 dimensional supersymmetric gauge theories the subject started with the work of [17, 18] and for the 5 dimensional cases in [31]. We give a short synopsis of these works focusing on the different geometries for the different Chern-Simons terms in the 5 dimensional case [31]. The main motivations are as follows.

Physically the higher genus terms calculable in the matrix model are only relevant if the theory is embedded in gravity, which according to our present understanding is consistently possible only by geometric engineering of the corresponding string geometries.

By moving in the moduli space, which is best described as the complex structure moduli space of the mirror geometry, one can connect the local matrix model descriptions of section 3 and 4 to each other and to the Chern-Simons type matrix models that have been studied in [22, 23, 24].

The 5-dimensional theories are labelled by different integer Chern-Simons terms $m \in \mathbb{Z}$, which make an important difference in the identification with the matrix model, discussed in section 5.5. For this reason and because the absence of world-sheet instantons makes the theories simpler to describe, we discuss mainly these cases [31].

The non-compact geometries for the 5-dimensional case are the same as the ones used

in the 4-dimensional geometric engineering after a limit [18]. We can therefore focus in section 3 on the discussion of this limit, which leads to 4-dimensional spectral curve $H(w, z; \mu_\alpha) = 0$. We explain why it does not depend on m .

For $m = 0$ and *ADE* gauge theories G there is another point in the moduli space where one extracts the partition function of Chern-Simons theory on the lens space S^3/D_G , where D_G is a discrete subgroup of $SU(2)$ acting on S^3 . The latter have a corresponding *ADE* classification and matrix model descriptions for these Chern-Simons theories have been suggested. The discussion in section 5.4 is necessary to specify the general Chern-Simons matrix model limit in [23].

5.1 Geometric engineering of 5-dimensional gauge theories

$\mathcal{N} = 1$ supersymmetric gauge theory in 5 dimensions can be geometrically engineered from M-theory on certain non-compact Calabi-Yau threefolds M . The basic geometry of M is that of a resolved *ADE* singularity fibered over $\mathbb{P}^1$. It gives rise to a $\mathcal{N} = 2$ supersymmetric gauge theory in five, and after a limit in four dimensions, whose gauge group G is the corresponding simply-laced group. The intersection of the $\text{rank}(G)$ $\mathbb{P}^1$ curves in the fiber is the negative Cartan matrix of G . For non-simply-laced groups the fibration has to have non-trivial monodromy, see [31]. Matter can be added by additional blow ups [18] leading to singular fibres [31] or by considering a higher genus base [34, 35].

The precise dictionary between manifolds M and the 5-dimensional gauge theories [31, 32, 33] contains a parameter $m \in \mathbb{Z}$, which labels a choice the Chern-Simons terms of the 5-dimensional gauge theory or equivalently the triple intersections of M . The choice $m = 0$ corresponds to the matrix model discussed in section 4. Since non-compact Calabi-Yau manifolds M on which M-theory compactifications give rise to 5-dimensional gauge theories with different Chern-Simons terms we need to describe how the differences in the non-compact mirror Calabi-Yau space W given by eq. (50) are reflected in the matrix model. At the end we propose matrix models describing topological string on a wide class of non-compact toric Calabi-Yau spaces, which correspond to the $SU(n)$ cases.

The different 5-dimensional theories, which however lead to the same gauge theory in 4 dimensions, can be specified by a particular choice of the normalized triple intersection numbers c_{ijk} of the non-compact Calabi-Yau, which appear in the 5-dimensional theory as Chern-Simons terms

$$\int c_{ijk} A_i \wedge F_j \wedge F_k . \quad (49)$$

Here the latin indices label the vector multiplets. Note that (49) is not gauge invariant, but it can be shown [36] that if M is spin with $p_1 = 0$, as it is the case if M is Calabi-Yau, the 5-dimensional integral over (49) shifts only by $2\pi\mathbb{Z}$. In M-theory on a Calabi-Yau M vector fields arise as $A_i = \int_{\mathcal{C}_i} C^{(3)}$, i.e. by reducing the three form field $C^{(3)}$ on a 2-cycle $\mathcal{C}_i$ in a class in $H_2(M)$. Similarly the field strength arises as $F_i = \int_{\mathcal{C}_i} (dC)^{(4)}$, so that (49) is just dimensional reduction of eleven dimensional Chern-Simons term $\int_M C^{(3)} \wedge (dC)^{(4)} \wedge (dC)^{(4)}$ with $c_{ijk} = \int_M \omega_i \wedge \omega_j \wedge \omega_k = D_i \cap D_j \cap D_k$, where ω_i are in $H^2(M)$ with compact support on $\mathcal{C}_i$ and D_i are dual divisors to $[\mathcal{C}_i]$. Both definitions of the triple intersection need a regularization on non-compact Calabi-Yau spaces. The intersections are largely determined by the Cartan matrix of G and in [31] it was shown that the discrete freedom in choosing different Chern-Simons terms is a discrete choice $m \in \mathbb{Z}$, which parametrizes that particular triple intersection mentioned above. For $SU(n)$ all triple intersections will be made explicit in the prepotential below.

5.2 Mirror symmetry for toric non-compact Calabi-Yau

Mirror symmetry implies that the data of the 5 dimensional gauge theory as well as of the topological string on M are captured by a Riemann surface (50) and a meromorphic differential λ . One crucial point is that spectral curve of the 5-dimensional matrix model discussed in the last section is identified with that Riemann surface. More precisely in the moduli space of the Riemann surface the partition function of the 5-dimensional gauge theory emerges as a holomorphic limit of the topological string partition function near the large radius point $\mu_\alpha = 0$. We review below how the mirror curve (50) for the geometry described in (5.1) is constructed.

The mirror manifold W of the manifold M is a conic bundle

$$H(w, z; \mu_\alpha) - uv = 0 \tag{50}$$

branched over $\mathbb{C}^* \times \mathbb{C}^*$ at a complex family of mirror curves $H(w, z; z_\alpha) = 0$, i.e. $u, v \in \mathbb{C}$, and $w, z \in \mathbb{C}^*$. With the geometry comes a canonical meromorphic differential

$$\lambda = \log(z) \frac{dw}{w} . \tag{51}$$

From our perspective this differential will determine the filling fractions of the matrix model.

For the $SU(n)$ cases the well known construction of non-compact toric mirror symmetry applies [18, 37], see [38] for a short review in the notation we use here. The manifolds M

and W are defined by charge vectors $Q_i^\alpha \in \mathbb{Z}$ of the toric group action. More precisely

$$M = (\mathbb{C}^{k+3} - Z)/(\mathbb{C}^*)^k, \quad (52)$$

where $(\mathbb{C}^*)^k$ acts by $x_i \mapsto \lambda_\alpha^{Q_i^\alpha} x_i$, $\alpha = 1, \dots, k$ on the complex coordinates x_i of $\mathbb{C}^{k+3}$ with $\lambda_\alpha \in \mathbb{C}^*$. The constraint $\sum_{i=1}^{k+3} Q_i^\alpha = 0$, $\forall \alpha$ is equivalent to $c_1(TM) = 0$. H is given by $H = \sum_{i=1}^{k+3} x_i$, where² $x_i \in \mathbb{C}^*$ are homogeneous coordinates w.r.t. an additional $\mathbb{C}^*$ -action and subject to the constraints

$$(-1)^{Q_0^\alpha} \prod_{i=1}^{k+3} x_i^{Q_i^\alpha} = \mu_\alpha, \quad \forall \alpha. \quad (53)$$

Here μ_α , $\alpha = 1, \dots, h_{11}^{comp}$, are complex moduli of (W, λ) , which are dual to the complexified Kähler parameters of M , while w and z in (50) denote the independent variables that remain after using the constraints (53) as well as the additional $\mathbb{C}^*$ -action on the x_i , mentioned above. $H(w, z, \mu_\alpha) = 0$ is central in the following discussion, because it is identified with the spectral curve of the matrix model.

5.3 Geometric engineering of 5-dimensional $SU(n)$ gauge theories

Let us now give the dependence of the charge vectors Q_k^α defining M and W on m . The simplest case arises for $SU(2)$ without matter. It was shown in [18] that the possible 5-dimensional geometry M corresponds in this case to the anticanonical line bundle over the Hirzebruch surfaces $\mathbb{F}_m = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(m))$, written here as the projectivization of line bundles over $\mathbb{P}^1$. The parameter m in this special case is precisely the $m \in \mathbb{Z}$ freedom to choose different Chern-Simons terms. Recall that this non-compact Calabi-Yau spaces over the Hirzebruch surfaces $\mathbb{F}_m$ are torically described by the 3-dimensional fan spanned by $\{n_1, n_2, \dots, n_5\} = \{(-1, 0, 1), (0, 0, 1)(1, 0, 1), (m, 1, 1), (1, -1, 1)\}$. The complete fan in the plane described by first two coordinates with the triangulation as shown in figure 4 defines torically the Hirzebruch surface $\mathbb{F}_m$.

Each point n_i corresponds to a divisor class $\hat{D}_i = \{x_i = 0\}$, where x_i are coordinates of the (52). These divisors are not independent, but fulfill the relations $\sum_{k=1}^r n_{k,i} \hat{D}_k$ for $r = 5$ and $i = 1, \dots, d = 3$. The triangulation specifies the intersection of M in the following way. $\hat{D}_i \hat{D}_j = 0$ if i and j are not part of one triangle. $\hat{D}_i \hat{D}_j \hat{D}_k = 1$ if n_i, n_j, n_k span a triangle in the triangulation. Restricted to the Hirzebruch surfaces one has $\sum_{k=1}^r n_{k,i} D_k$

²We use the same symbol x_i for the different coordinates of M and W .




Figure 4: Toric fans for $\mathbb{F}_m$.

for $r = 5$ and $i = 1, \dots, d = 2$, $D_i \cap D_j = 1$, $i, j > 1$ if n_i, n_j are on a line³. There are two independent classes in $\mathbb{F}_m$, $F = D_4 = D_5$ and $S = D_1 = S' - mF$ with $S' = D_3$. With the above information and the ring structure of the intersections it is easy to calculate all toric intersections. For the Hirzebruch surface $FS = 1$, $F^2 = 0$, $S'S = 0$ and $S^2 = -2$. For M there is now a 2-dimensional complex dimensional compact divisor $H = D_1$, which represents the Hirzebruch surface itself. $H^3 = 4$, $H\hat{S}^2 = -m$, $H\hat{F}^2 = 0$ and $H\hat{F}\hat{S} = 1$, where $\hat{S}$ and $\hat{S}$ are non-compact divisors of M . Note that the triple intersections among the non-compact divisors corresponding to points on the boundary of the fan are not well defined.

The charge vectors are coefficients of linear relations among the points $\sum_i Q_i^\alpha n_i = 0$. The one which do actually span the Mori cone are calculated using the secondary polytop to be $Q^1 = (1, -2, 1, 0, 0)$ and $Q^2 = (0, m - 2, -m, 1, 1)$. They correspond to curve classes and the intersection number of the α 'th curve with the divisor D_i is encoded in Q_i^α . The Kähler cone is dual to the Mori cone and it is easy to see that it is spanned by F and $H = S + (m + 1)F$.

These geometries are easily generalized to $SU(n)$. To distinguish n odd and n even we define $\Delta(n) = \frac{1}{2}(1 - (-1)^n)$. The points of the 3-dimensional fan and the Mori cone are

³We use hatted characters for non-compact divisors of M that restrict to compact divisors on $\mathbb{F}_m$.

given, with $n_l = \frac{\Delta(n)-n}{2}$ and $n_u = \frac{n+\Delta(n)}{2}$, as

n_i	Q^1	Q^2	Q^3	$\dots$							Q^n
$(n_l, 0, 1)$	1	0	0	$\dots$	0	0	0	$\dots$	0	0	0
$(n_l + 1, 0, 1)$	-2	1	0	$\dots$	0	0	0	$\dots$	0	0	0
$(n_l + 2, 0, 1)$	1	-2	1	$\dots$	0	0	0	$\dots$	0	0	0
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\dots$	$\vdots$	$\vdots$	$\vdots$	$\dots$	$\vdots$	$\vdots$	$\vdots$
$(0, 0, 1)$	0	0	0	$\dots$	1	-2	1	$\dots$	0	0	$(m - 2 + \Delta(n))$
$(1, 0, 1)$	0	0	0	$\dots$	0	1	-2	$\dots$	0	0	$-m - \Delta(n)$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\dots$	$\vdots$	$\vdots$	$\vdots$	$\dots$	$\vdots$	$\vdots$	$\vdots$
$(n_u - 2, 0, 1)$	0	0	0	$\dots$	0	0	0	$\dots$	1	-2	1
$(n_u - 1, 0, 1)$	0	0	0	$\dots$	0	0	0	$\dots$	0	1	-2
$(n_u, 0, 1)$	0	0	0	$\dots$	0	0	0	$\dots$	0	0	1
$(m, 1, 1)$	0	0	0	$\dots$	0	0	0	$\dots$	0	0	0
$(\Delta(n), -1, 1)$	0	0	0	$\dots$	0	0	0	$\dots$	0	0	0

n odd



n even




Figure 5: Toric fans for the non-compact toric Calabi-Yau corresponding to $SU(n)$ theory with the Chern-Simons term parameterized by m .

One can see from the pictures that the inner points in the diagram correspond to toric compact divisors of the type $\mathbb{F}_k$ in the non-compact threefold, which can be easily seen to be Hirzebruch surfaces. We denote these divisors H_i , $i = 1, \dots, N - 1$. From the left to the right $H_i = \mathbb{F}_{|m+2i-n|}$, $i = 1, \dots, n - 1$. With the description for the calculation of the intersection of the H_i it is easy to confirm [31] that the cubic prepotential, which is not corrected by instantons for 5-dimensional theories is $F = \frac{1}{2} \sum_{i < j} |a_i - a_j|^3 + m \sum_i a_i^3 = \sum_{i_1, i_2, i_3} H_{i_1} H_{i_2} H_{i_3} \prod_{r=1}^3 (a_{i_r+1} + \dots + a_{i_n})$.

The standard parametrization of mirror symmetry using the Mori cone in (53) implies that $\mu_\alpha = 0$ is a point of maximal unipotent monodromy and that near this point $t_\alpha \sim \frac{1}{2\pi i} \log(\mu_\alpha)$, with t_α the complexified volumes of the curves corresponding to the Mori cone. For $\alpha = 1, \dots, n-1$ these curves are the fibres of the Hirzebruch surfaces. Geometrically their size is proportional to the mass of the W bosons in string units M_{string} . It is given by the simple roots, i.e. $t_\alpha = a_\alpha - a_{\alpha+1}$, $\alpha = 1, \dots, n-1$. The parameter $t_b := t_n$ corresponds to the complexified size of the base $\mathbb{P}^1$. The size appears as imaginary part while the B -field integral over the curve is the real part.

5.4 4-dimensional Seiberg-Witten limit

We come now to the engineering of the 4-dimensional gauge theory. The aim is to derive from the 5-dimensional spectral curve $H(w, z; \mu_\alpha) = 0$ in a limit the 4-dimensional Seiberg-Witten (7) curve and show that λ reduces to dS . Moreover in the 4-dimensional limit the differences given by the parameter m should vanish. While the 5-dimensional gauge theory discussed above is recovered in the large radius limit, the construction of the 4-dimensional gauge theory requires a double scaling limit in a region of the moduli space, called Seiberg-Witten point, where world-sheet instantons dominate. In fact it has been made explicit in [18] how the individual gauge theory instantons contributions of Seiberg-Witten gauge theory are encoded in a resummation of the world-sheet instantons.

Non-abelian gauge groups in string theory have been first understood in the heterotic construction. For finding the Seiberg-Witten point it is instructive to recall the duality between the heterotic and type II strings. It states [17] that the heterotic dilaton S is geometrized and identified with $\frac{1}{4\pi i} t_b$, i.e. the size of the base of the fibration discussed in 5.3. In the heterotic string one has $\text{Re}(S) = \frac{1}{g^2}$ so that $\text{Im}(t_b) = \frac{4\pi}{g^2}$. The asymptotic free $g \rightarrow 0$ region of Seiberg-Witten must be at large volume of the base $\mathbb{P}^1$ and hence near $\mu_b := \mu_n = 0$. However a double scaling limit, parametrized below by $\epsilon \rightarrow 0$ must be taken to keep the mass of the W bosons finite, while taking the string scale to infinity. The mass of the W bosons is the only scale and the one-loop running of the gauge coupling in $\mathcal{N} = 2$ gauge theory implies $\frac{1}{g^2} \sim \log\left(\frac{M_W}{\Lambda}\right)$. Since W -boson comes from branes wrapping the fibers one has $t_b \sim C \log(t_f)$, where t_f refers to generic volumes of the fiber. The constant C can be determined from the dependence of the instanton contribution with instanton number k namely $\exp\left(-\frac{k}{g^2}\right) \sim \left(\frac{1}{t_f^{2n}}\right)^k$, where $\hat{t}_f$ refers volumes of the fibers, i.e. masses of the W bosons, in gauge theory units. Putting the above facts together, the

double scaling limit is $\exp(t_b) = (\epsilon\Lambda)^{2n}$ and $t_f = \epsilon\hat{t}_f$. In the B-model geometry, i.e. the manifold W , t_f is mapped to period integrals over S^1 , which come from periods over S^3 in the compact geometry. That is the Seiberg-Witten curve is near point where the divisor $\mu_n = 0$ intersects $n - 1$ conifold divisors in the moduli space. The double scaling limit can also be viewed as resolving the point of tangency between the divisors of this intersection point, to obtain normal crossing divisors.

Let us see for the $SU(2)$ cases how (7) and dS is recovered in the limit from the $5d$ curve. According to the table in section 5.3 and the description how to obtain $H(w, z; \mu_\alpha)$ from (53) we get

$$H(w, z; \mu_\alpha) = w + \frac{\mu_f z^{2-m}}{w} + 1 + z + \mu_b z^2 = 0 \quad (54)$$

as the defining equation of the 5-dimensional curve. Here we identified in $H = \sum_{i=1}^5 x_i$, $w := x_5$ and $z := x_2$. We set $x_3 = 1$ by the $\mathbb{C}^*$ -action and eliminated x_4 and x_1 by (53). Now we calculate the conifold discriminant and find that it behaves like $\Delta = (1 - 4\mu_f)^2 + \mathcal{O}(\mu_b)$, where all the m -dependence is in higher order in μ_b . M_W^2 is proportional to Δ which and the scaling argument implies that the finite gauge theory coordinate $u \sim \hat{t}_f^2$ should appear as

$$\begin{aligned} \eta_1 &= (1 - 4\mu_f) = \epsilon^2 u, \\ \eta_2 &= \frac{\sqrt{\mu_b}}{(1 - 4\mu_f)} = \frac{\Lambda^2 e^{-S_0/2}}{u}, \end{aligned} \quad (55)$$

here $e^{-S_0/2}$ is an a priori undetermined constant and (55) are just the blow up variables to resolve the tangency between $\mu_b = 0$ and $\Delta = 0$. An important point is that z is a $\mathbb{C}^*$ variable in $H(w, z; \mu_\alpha) = 0$, but in (7) x is a $\mathbb{C}$ variable, which is clearly reflected by the $\log(z)$ dependence of λ as opposed to the x dependence of dS . So we can think of (7) as a compactification of $H = 0$ in a special limit of its parameter space. To match the the differentials we must identify in leading order $z = 1 - \epsilon\tilde{x}$. Rescaling $\tilde{x} = \frac{\hat{x}}{\epsilon\sqrt{a_f}}$ and eliminating the $\tilde{x}$ term by $\tilde{x} = x + \frac{1}{2} \left(\frac{1}{\sqrt{\mu_f}} + 2\sqrt{\mu_f} \right)$ one gets independent of m in leading order $w + \frac{\sqrt{\mu_b}}{w} = -x^2 - \left(1 - \frac{1}{4\mu_f} \right)$. Rescaling further $w \rightarrow w\sqrt{\mu_b}$ and $x \rightarrow \frac{i(\mu_b)^{\frac{1}{4}}}{\Lambda}x$ and using (55) we get (7).

5.5 Matrix model description of the $m \neq 0$ cases

For $m \neq 0$ the microscopic matrix model can be obtained starting with the modified Nekrasov partition function [33], which is written as in (12) with a linear modification and

quadratic modification in λ by the second Casimir operator

$$\kappa_\lambda = \sum_i \lambda_i(\lambda_i + 1 - 2i) \quad (56)$$

and a linear term in λ as

$$Z_{5d}^{SU(n)} = \sum_{\vec{\lambda}=(\lambda^{(1)}, \dots, \lambda^{(n)})} (\beta\Lambda)^{2n|\vec{\lambda}|} e^{-m\beta \sum_l |\lambda^{(l)}| a_l + \hbar \kappa_{\lambda^{(l)}}} Z_{\beta, \vec{\lambda}}. \quad (57)$$

It is straightforward to check [33], that this expression is reproduced by the applications of the topological vertex [39] to the geometries discussed in section 5.3.

Likewise it is simple to see, that if we apply the same steps as in section 4 to $Z_{\beta, \vec{\lambda}}$ we get (39), except that we cannot view the whole expression, which follows from (57) as a matrix integral, because of the different linear coupling (distinguished by a_l) to n sets of eigenvalues in the potential. These linear terms are the only relevant change in the potential, since the shifts in the definition (32) give just rise to an overall constant and moreover the quadratic terms from the second Casimir operator will vanish in the $\hbar \rightarrow 0$ limit.

In order to convert this to matrix model one has to introduce n matrices. The procedure is similar as in [23]. A linear term of the form $a_l \sum_i u_i^{(l)}$ in a quadratic potential, which is integrated against the deformed measure $\Delta_q(u^{(l)})$ (38) can be absorbed into an interaction between the different groups of eigenvalues $\Delta_q(u^{(l)}, u^{(k)}) = \prod_{i,j} 2 \sinh \left(\frac{u_i^{(l)} - u_j^{(k)}}{2} \right)$ in the measure. This holds for the potential (44). After viewing the n groups of $u^{(l)}$, $l = 1, \dots, n$ as the eigenvalues of n matrices the interacting factors in the measure can be exponentiated to give a relatively complicated interaction term between the matrices [23]. The matrix models for $m \neq 0$ describe, e.g. for $Su(2)$ and $m = 1$ the anticanonical bundle over $\mathbb{F}_1$, which by a blow down becomes $\mathbb{P}^2$. So it seems possible to write down a microscopic multimatrix model, which reproduces in a limit the simplest non-compact toric Calabi-Yau $\mathcal{O}(-3) \rightarrow \mathbb{P}^2$.

6 Relations to other matrix models

In this section we show that various well-known and seemingly unrelated matrix models arise in various limits of the 5-dimensional matrix model derived in section 4. These interesting cases include Chern-Simons matrix models for lens spaces, as well as matrix models for line bundles over $\mathbb{P}^1$. Moreover, in the 4-dimensional limit the 5-dimensional

model reduces to the matrix model from section 3, which, as we discuss below, can be regarded as a generalization of the Eguchi-Yang matrix model (which itself corresponds just to the $n = 1$ case). In a sense, in this way we exhibit a part of a vast and unified matrix model landscape, whose various limits were introduced and considered in the past in an unrelated and independent fashion. We discuss these limiting cases in the following subsections.

6.1 Chern-Simons matrix model for lens spaces

The orbifold limit in the $m = 0$ case is at the point in the moduli space where all t_i , $i = 1, \dots, n$, are small. In the complex structure variables this is near the vanishing of $w_i = 1 - \frac{z_i}{z_{i+1}}$, $i = 1, \dots, n-1$ and $w_n = \frac{1}{\sqrt{z_{n-1}} \left(1 - \frac{z_n}{z_{n-1}}\right)}$. Here is another matrix model known describing this topological string theory in the orbifold point [23]. Its spectral curve was analyzed in [24, 25] and claimed to be the curve $H(x, y; z) = 0$ obtained from (53) and the table in section 5.3, as described above.

The matrix model presented in [23] looks almost the same as ours – the only difference is that its potential contains only the quadratic term. It does not contain neither the linear term in t , nor the dilogarithms. The absence of the linear term is easy to explain, as $t = 0$ should in principle correspond to the orbifold limit. We also note that the dilogarithms are suppressed when their arguments have large negative values. It would be interesting to understand the relation between a continuation from the large radius to the orbifold point in more detail. We also note that slightly more general models for orbifold points corresponding to arbitrary lens spaces $L(p, q)$ we recently found in [41]. They also have just the quadratic potential and the deformed measure, so belong to the same family of models as those described above.

6.2 Matrix models for line bundles over $\mathbb{P}^1$

In [21] topological strings for Calabi-Yau manifolds of the form

$$X_p = \mathcal{O}(p-2) \oplus \mathcal{O}(-p) \rightarrow \mathbb{P}^1$$

were considered. Partition functions for such theories can be computed using the topological vertex [39]

$$Z_{X_p} = \sum_{\lambda} C_{\lambda \bullet \bullet} C_{\lambda^t \bullet \bullet} q^{(p-1)\kappa_{\lambda}/2} (-1)^{|\lambda|} e^{-t|\lambda|},$$

where the sum is performed over all two-dimensional partitions λ , $\kappa_\lambda = \sum_i \lambda_i(\lambda_i - 2i + 1)$ is the second Casimir, $q = e^{-gs}$, and t is the Kähler parameter. The vertex amplitude in this case $C_{R\bullet\bullet} = P_q(\lambda)^{1/2}$ is a square root of the deformed Plancherel measure (5), so it leads to the same partition function as considered in [14]. The case $p = 1$ can be considered as the 5-dimensional $U(1)$ theory, which is a special case $n = 1$ of our general result. It was shown in [21] that the above partition function can be written as a one-cut matrix model with a deformed measure and the potential

$$V(u) = (t - (p - 1))u + \frac{p}{2}u^2 + 2 \text{Li}_2(e^u).$$

For $p = 1$ this is indeed the potential which we obtain in $n = 1$ case. For arbitrary p only the linear and quadratic terms are modified (the latter due to the Casimir κ_λ in Z_{X_p}).

6.3 Eguchi-Yang matrix model

We also note that in the limit of the 4-dimensional theory and for $n = 1$ our matrix model with the potential (26) is closely related to the $\mathbb{CP}^1$ matrix model introduced by Eguchi and Yang

$$\begin{aligned} Z^{EY} &= \int \mathcal{D}M e^{-2N \text{Tr } V^{EY}(M)}, \\ V^{EY} &= (-M + M \log M) - \sum_{i=1}^{\infty} t_{i-1,Q} M^i - \sum_{i=1}^{\infty} t_{i,P} M^i (\log M - \sum_{j=1}^i \frac{1}{j}). \end{aligned}$$

This matrix model was introduced in [19, 20], where it was also shown that its partition function Z^{EY} reproduces the partition function of the topological $\mathbb{CP}^1$ model considered in [40], i.e. the topological sigma-model coupled to the two-dimensional gravity. This model has two observables denoted $P \equiv \sigma_0(P)$ and $Q \equiv \sigma_0(Q)$, which correspond respectively to the identity and the Kähler class of $\mathbb{P}^1$. In addition, two infinite series of descendants $\sigma_i(P)$ and $\sigma_i(Q)$, for $i = 1, 2, \dots$, arise in this theory upon coupling to gravity. Coupling constants corresponding to these descendants are denoted by $t_{i,P}$ and $t_{i,Q}$, and they appear in the Eguchi-Model as coefficients in two infinite series in the potential.

We immediately see that a special case of the Eguchi-Yang model corresponding to $t_{i,P} = t_{i,Q} = 0$ for $i \geq 1$ is identical with the $n = 1$ case of our matrix model (26), with our t identified with the coupling $t_{0,Q}$ in V^{EY} . From physical point of view the vanishing of all coupling constants corresponds to replacing a matrix M by a chiral superfield [42]. From the model with $n = 1$ and vanishing coupling constants one can derive the limit

shape of partitions given by the arcsin-law [14, 43]. Our matrix model for Seiberg-Witten theory (26) is therefore a generalization of the Eguchi-Yang matrix model to arbitrary n . More generally, setting all $t_{i,P} = 0$ but taking arbitrary $t_{i,Q}$ is identical with the matrix model considered in [14], with arbitrary Casimirs taken into account. We note that some subtleties, advantages and drawbacks of the comparison of the sums over partitions arising from gauge theory instanton calculus, to the matrix model of the Eguchi-Yang type, are also discussed in [43].

7 Conclusions and discussion

In this paper we provided a formulation of Seiberg-Witten theory in terms of matrix models. For 4-dimensional theories we derived 1-matrix models with a standard measure containing the Vandermonde determinant, while the 1-matrix models for 5-dimensional theories have a deformed measure. We also extended this approach to a matrix model formulation of topological string theory on Calabi-Yau manifolds which are related to Seiberg-Witten theories by geometric engineering. In general this involves multi-matrix models.

The 1-matrix models which we propose were derived in two steps. Firstly, we introduced in the Nekrasov partition function an auxiliary parameter N , and rewrote this partition function as a matrix integral over matrices of size N . The matrix models obtained this way are rather non-standard 1-matrix models, with potentials which contain such special functions as logarithms of Γ functions. Then, the partition functions of the Seiberg-Witten theory were recovered in the 't Hooft limit. In this limit various simplifications take place and we end up with matrix potentials containing logarithmic and dilogarithmic terms.

The derivation of these matrix models is just an initial step, which should be followed by a detailed analysis of their properties. We use the Migdal-Muskhelishvili formula for the resolvent to check that the 1-matrix models for the 5-dimensional theories yield the expected spectral curves with the correct differential λ defining the filling fractions. The 4-dimensional spectral curves and the corresponding differential dS are reproduced by in the gauge theory limit of geometric engineering. It could have also been derived with the Migdal-Muskhelishvili formalism for the resolvent in 4-dimensional theory directly.

The general framework for solving matrix models based just on the underlying spectral curve and the differential, which was developed by Eynard and Orantin [11, 44] and checked in very similar situations in [45, 46], should give the higher genus, as well as open amplitudes

in all our cases⁴. The holomorphic anomaly equations and the gap condition applies likewise to the 4-dimensional cases [12] and to the 5-dimensional models [38]. It provides the most efficient way to calculate the closed higher genus amplitudes.

Certainly a perturbative expansion of the matrix models discussed here is possible. Of course we hope that the knowledge of the microscopic matrix model action will provide a route towards better understanding non-perturbative effects in Seiberg-Witten theory and topological strings. Techniques of analyzing non-perturbative effects in matrix models, being currently developed by [47, 48], require the knowledge of the potential, which we provide in this paper. They would also be of great interest from the point of view of black hole physics and OSV conjecture [49], and complementary to the proposition discussed in [50, 51].

An important aspect of the matrix models derived here is their relation to other families of matrix models. Firstly, matrix models for the $SU(n)$ 4-dimensional Seiberg-Witten theory can be regarded as a generalization of the the Eguchi-Yang matrix model (which corresponds to $n = 1$). Higher Casimir operators, which we did not analyse in this paper, but in the context of $n = 1$ model they were taken into account into [14, 43], would correspond to gravitational descendants in the Eguchi-Yang model. In the same spirit, matrix models for 5-dimensional $SU(n)$ theory derived here are generalizations of the matrix model for bundles over $\mathbb{P}^1$ considered in [21] (which also corresponds to $n = 1$). It is also reassuring that the models which we derived contain no other terms than logarithms and dilogarithms, which also appeared in [21]. While our models, as well as those in [21], correspond to the large radius point in the moduli space, it is also possible to consider other regions of the moduli space. In particular, in the limit where dilogarithms in the potential for the 5-dimensional theory are suppressed, we recovered Chern-Simons matrix models for lens spaces which correspond to orbifold points in the moduli space [23, 24, 25]. The similarity of the microscopic description at the large radius point and at the orbifold point (just the suppression of dilogarithmic terms) is surprising in view of highly involved transformations of partition functions between these points.

The above results indicate a possible unified microscopic description of the multitude of matrix models, which describe various regions in the moduli space. Similar ideas of the unification of matrix models to some overall matrix-model-M-theory (in analogy to string theory) were advocated in [52, 53].

Our results should also provide a direct link between fermionic formulations of matrix

⁴Some checks in 4-dimensional Seiberg-Witten theory have been made in [13].

models, gauge theories and topological strings. The fermionic viewpoint on matrix models, intimately related to conformal field theory techniques, was originally advanced by Kostov [54]. Recently the (matrix model) formalism of Eynard and Orantin was also rephrased in a fermionic language [55]. On the other hand, topological strings and gauge theories can be formulated in terms of two-dimensional chiral fermions too [10, 39, 56, 57, 58, 59]. Such fermions live on curves which, in the present context, are simultaneously matrix model spectral curves and Seiberg-Witten curves. This convinces that all those fermions have a common origin. Moreover, in [57] a formalism was presented which relates both matrix models and Seiberg-Witten theory to $\mathcal{D}$ -modules. Finding how matrix models presented in this paper fit into this formalism would certainly be of interest.

Let us also stress that the idea of expressing gauge theory quantities in terms of matrix integrals has been explored to much extent in the context of $\mathcal{N} = 1$ theories within the so-called Dijkgraaf-Vafa formalism [60, 61]. $\mathcal{N} = 1$ theories can be treated as deformations of $\mathcal{N} = 2$ theories, and it was also shown that certain scaling of Dijkgraaf-Vafa matrix models leads to the $\mathcal{N} = 2$ answers [61, 62]. The difference with the case considered in this paper is that now we get the answer which gives $\mathcal{N} = 2$ results without a need for any rescaling. Some other matrix models or related ideas corresponding to gauge theories were also proposed in [63, 64, 65, 66, 67]. It would be interesting to relate these points of view.

Acknowledgments

We would like to thank Rainald Flume for many comments, suggestions and collaboration in the initial stages of this project. We also appreciate useful suggestions and discussions with Vincent Bouchard, Marcos Mariño, Andrei Marshakov, Andrei Mironov, Nikita Nekrasov and Cumrun Vafa. P.S. would like to thank the organizers of the 6th Simons Workshop in Mathematics and Physics in Stony Brook, where parts of this project were done, for hospitality and inspiring atmosphere. The research of P.S. was supported by the Humboldt Fellowship.

Appendix - elliptic integrals

To derive the spectral curve for 5-dimensional $SU(2)$ theory we need to perform the following integral

$$\int^v \frac{v^{-1}(z-v)^{-1} dv}{\sqrt{(v-a)(v-b)(v-c)(v-d)}} = g_1 F(y(v), k) + g_2 \Pi(y(v), \delta(0), k) + g_3 \Pi(y(v), \delta(z), k),$$

which is expressed as a combination of the elliptic functions, with the following notation. The coefficients in this combination are equal

$$g_1 = \frac{2}{a(a-z)\sqrt{(b-c)(a-d)}}, \quad g_2 = \frac{2(b-a)}{zab\sqrt{(b-c)(a-d)}}, \quad (58)$$

$$g_3 = \frac{2(b-a)}{z(z-a)(z-b)\sqrt{(b-c)(a-d)}},$$

while the arguments of F and Π are given as

$$y(v) = \frac{(a-d)(b-v)}{(b-d)(a-v)}, \quad \delta(z) = \frac{1}{y(z)}, \quad k^2 = \frac{1}{y(c)} = \frac{(b-d)(a-c)}{(a-d)(b-c)}. \quad (59)$$

The elliptic functions F and Π , of the first and the third kind respectively, are

$$F(y, k) = \int_0^y \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}, \quad \Pi(y, \delta, k) = \int_0^y \frac{dt}{(1-\delta t^2)\sqrt{(1-t^2)(1-k^2t^2)}}. \quad (60)$$

The complete elliptic integrals are defined as $K(k) \equiv F(1, k)$ and $\Pi(\delta, k) \equiv \Pi(1, \delta, k)$. Of particular importance in our computation are the following addition formulas [68]

$$F(\sin \vartheta, k) \pm F(\sin \beta, k) = F(\sin \varphi, k), \quad (61)$$

$$\Pi(\sin \vartheta, \delta, k) \pm \Pi(\sin \beta, \delta, k) = \Pi(\sin \varphi, \delta, k) \pm \sqrt{\frac{\delta}{(\delta-1)(\delta-k^2)}} \operatorname{artanh} \Theta, \quad (62)$$

where

$$\Theta = \frac{\sin \vartheta \sin \beta \sin \varphi \sqrt{\delta(\delta-1)(\delta-k^2)}}{1 - \delta \sin^2 \varphi + \delta \sin \vartheta \sin \beta \cos \varphi \sqrt{1 - k^2 \sin^2 \varphi}}, \quad (63)$$

and the angle φ is determined as

$$\cos \varphi = \frac{\cos \vartheta \cos \beta \mp \sin \vartheta \sin \beta \sqrt{(1 - k^2 \sin^2 \vartheta)(1 - k^2 \sin^2 \beta)}}{1 - k^2 \sin^2 \vartheta \sin^2 \beta}. \quad (64)$$

References

- [1] N. Seiberg, E. Witten, *Monopole Condensation, And Confinement In $N=2$ Supersymmetric Yang-Mills Theory*, Nucl. Phys. B426 (1994) 19-52 [hep-th/9407087].
- [2] N. Seiberg, E. Witten, *Monopoles, Duality and Chiral Symmetry Breaking in $N=2$ Supersymmetric QCD*, Nucl. Phys. B431 (1994) 484-550 [hep-th/9408099].
- [3] A. Bilal, *Duality in $N=2$ SUSY $SU(2)$ Yang-Mills Theory: A pedagogical introduction to the work of Seiberg and Witten*, hep-th/9601007.
- [4] A. Klemm, *On the Geometry behind $N=2$ Supersymmetric Effective Actions in Four Dimensions*, hep-th/9705131.
- [5] E. D'Hoker, D. Phong, *Lectures on Supersymmetric Yang-Mills Theory and Integrable Systems*, hep-th/9912271.
- [6] N. Dorey, T. Hollowood, V. Khoze, M. Mattis, *The Calculus of Many Instantons*, Phys. Rept. 371 (2002) 231-459 [hep-th/0206063].
- [7] K. Iga, *What do Topologists want from Seiberg–Witten theory? (A review of four-dimensional topology for physicists)*, Int. J. Mod. Phys. A17 (2002) 4463-4514 [hep-th/0207271].
- [8] J. Labastida, M. Marino, *Topological Quantum Field Theory and Four Manifolds* Math. Phys. Studies 25.
- [9] N. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, Adv. Theor. Math. Phys. 7 (2004) 831-864 [hep-th/0206161].
- [10] N. Nekrasov, A. Okounkov, *Seiberg-Witten theory and random partitions*, hep-th/0306238.
- [11] B. Eynard, N. Orantin, *Invariants of algebraic curves and topological expansion*, math-ph/0702045.
- [12] M. x. Huang, A. Klemm, *Holomorphic anomaly in gauge theories and matrix models*, JHEP 0709 (2007) 054 [hep-th/0605195].
- [13] M. x. Huang, A. Klemm, *Modularity versus holomophicity in gauge theories and matrix models*, JHEP **0709**, 054 (2007) [hep-th/0605195].

- [14] B. Eynard, *All order asymptotic expansion of large partitions*, 0804.0381 [math-ph].
- [15] A. Vershik, S. Kerov, *Asymptotics of the Plancherel measure of the symmetric group and the limit form of Young tableaux*, Soviet Math. Dokl. 18 (1977) 527-531.
- [16] B. Logan, L. Shepp, *A variational problem for random Young tableaux*, Adv. Math. 26 (1977) 206-222.
- [17] S. Kachru, A. Klemm, W. Lerche, P. Mayr, C. Vafa, *Nonperturbative results on the point particle limit of $N=2$ heterotic string compactifications*, Nucl. Phys. B459 (1996) 537 [hep-th/9508155].
- [18] S. Katz, A. Klemm, C. Vafa, *Geometric engineering of quantum field theories*, Nucl. Phys. B497 (1997) 173-195 [hep-th/9609239].
- [19] T. Eguchi, S. Yang, *The Topological $\mathbb{CP}^1$ Model and the Large- N Matrix Integral* Mod. Phys. Lett. A9 (1994) 2893-2902 [hep-th/9407134].
- [20] T. Eguchi, K. Hori, S. Yang, *Topological σ -Models and Large- N Matrix Integral*, Int. J. Mod. Phys. A10 (1995) 4203 [hep-th/9503017].
- [21] N. Caporaso, L. Griguolo, M. Marino, S. Pasquetti, D. Seminara, *Phase transitions, double-scaling limit, and topological strings*, Phys. Rev. D75 (2007) 046004 [hep-th/0606120].
- [22] M. Marino, *Chern-Simons theory, matrix integrals, and perturbative three-manifold invariants*, Commun. Math. Phys. 253 (2004) 25-49 [hep-th/0207096].
- [23] M. Aganagic, A. Klemm, M. Marino, C. Vafa, *Matrix Model as a Mirror of Chern-Simons Theory*, JHEP 0402 (2004) 010 [hep-th/0211098].
- [24] N. Halmagyi, V. Yasnov, *The Spectral Curve of the Lens Space Matrix Model*, hep-th/0311117.
- [25] N. Halmagyi, T. Okuda, V. Yasnov, *Large N Duality, Lens Spaces and the Chern-Simons Matrix Model*, JHEP 0404 (2004) 014 [hep-th/0312145].
- [26] H. Nakajima, K. Yoshioka, *Instanton counting on blowup. I. 4-dimensional pure gauge theory*, math.AG/0306198.

- [27] H. Nakajima, K. Yoshioka, *Instanton counting on blowup. II. K-theoretic partition function*, math.AG/0505553.
- [28] N. Nekrasov, S. Shadchin, *ABCD of instantons*, Commun. Math. Phys. 252 (2004) 359-391 [hep-th/0404225].
- [29] A. Migdal, *Loop Equations and $1/N$ Expansion*, Phys. Rept. 102 (1983) 199-290.
- [30] L. Chekhov, A. Marshakov, A. Mironov, D. Vasiliev, *Complex Geometry of Matrix Models*, hep-th/0506075.
- [31] K. A. Intriligator, D. R. Morrison, N. Seiberg, *Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau spaces*, Nucl. Phys. B497 (1997) 56 [hep-th/9702198].
- [32] A. Iqbal, A. Kashani-Poor, *$SU(N)$ Geometries and Topological String Amplitudes*, Adv. Theor. Math. Phys. 10 (2006) 1-32 [hep-th/0306032].
- [33] Y. Tachikawa, *Five-dimensional Chern-Simons terms and Nekrasov's instanton counting*, JHEP 0402 (2004) 050 [hep-th/0401184].
- [34] S. H. Katz, D. R. Morrison, M. Ronen Plesser, *Enhanced Gauge Symmetry in Type II String Theory*, Nucl. Phys. B477 (1996) 105 [hep-th/9601108].
- [35] A. Klemm, P. Mayr, *Strong Coupling Singularities and Non-abelian Gauge Symmetries in $N = 2$ String Theory*, Nucl. Phys. B469 (1996) 37 [hep-th/9601014].
- [36] E. Witten, *Phase Transitions In M-Theory And F-Theory*, Nucl. Phys. B471 (1996) 195 [hep-th/9603150].
- [37] K. Hori, C. Vafa, *Mirror symmetry*, hep-th/0002222.
- [38] B. Haghighat, A. Klemm, M. Rauch, "Integrability of the holomorphic anomaly equations," 0809.1674 [hep-th].
- [39] M. Aganagic, A. Klemm, M. Marino, C. Vafa, *The topological vertex*, Commun. Math. Phys. 254 (2005) 425-478 [hep-th/0305132].
- [40] R. Dijkgraaf, E. Witten, *Mean Field Theory, Topological Field Theory, And Multimatrix Models*, Nucl. Phys. B342 (1990) 486-522.

- [41] A. Brini, L. Griguolo, D. Seminara, A. Tanzini, *Chern-Simons theory on $L(p, q)$ lens spaces and Gopakumar-Vafa duality*, 0809.1610 [hep-th].
- [42] S. Cecotti, C. Vafa, *On classification of $N=2$ supersymmetric theories*, Commun. Math. Phys. 158 (1993) 569-644 [hep-th/9211097].
- [43] A. Marshakov, N. Nekrasov, *Extended Seiberg-Witten Theory and Integrable Hierarchy*, JHEP 01 (2007) 104 [hep-th/0612019].
- [44] B. Eynard, *All genus correlation functions for the hermitian 1-matrix model*, JHEP 0411 (2004) 031 [hep-th/0407261].
- [45] M. Marino, *Open string amplitudes and large order behavior in topological string theory*, hep-th/0612127.
- [46] V. Bouchard, A. Klemm, M. Marino, S. Pasquetti, *Remodeling the B-model*, 0709.1453 [hep-th].
- [47] M. Marino, R. Schiappa, M. Weiss, *Nonperturbative Effects and the Large-Order Behavior of Matrix Models and Topological Strings*, 0711.1954 [hep-th].
- [48] M. Marino, *Nonperturbative effects and nonperturbative definitions in matrix models and topological strings*, 0805.3033 [hep-th].
- [49] H. Ooguri, A. Strominger, C. Vafa, *Black Hole Attractors and the Topological String*, Phys.Rev. D70 (2004) 106007 [hep-th/0405146].
- [50] M. Aganagic, H. Ooguri, N. Saulina, C. Vafa, *Black Holes, q -Deformed 2d Yang-Mills, and Non-perturbative Topological Strings*, Nucl. Phys. B715 (2005) 304-348 [hep-th/0411280].
- [51] M. Aganagic, D. Jafferis, N. Saulina, *Branes, Black Holes and Topological Strings on Toric Calabi-Yau Manifolds*, JHEP 0612 (2006) 018 [hep-th/0512245].
- [52] A. Alexandrov, A. Mironov, A. Morozov, *M-Theory of Matrix Models*, hep-th/0605171.
- [53] A. Alexandrov, A. Mironov, A. Morozov, *Instantons and Merons in Matrix Models*, hep-th/0608228.
- [54] I. Kostov, *Conformal Field Theory Techniques in Random Matrix models*, hep-th/9907060.

- [55] R. Dijkgraaf, C. Vafa, *Two Dimensional Kodaira-Spencer Theory and Three Dimensional Chern-Simons Gravity*, 0711.1932 [hep-th].
- [56] R. Dijkgraaf, L. Hollands, P. Sułkowski, C. Vafa, *Supersymmetric gauge theories, intersecting branes and free fermions*, JHEP 0802 (2008) 106 [0709.4446 [hep-th]].
- [57] R. Dijkgraaf, L. Hollands, P. Sułkowski, *Quantum Curves and $\mathcal{D}$ -modules*, 0810.4157 [hep-th].
- [58] P. Sułkowski, *Calabi-Yau crystals in topological string theory*, PhD thesis [0712.2173 [hep-th]].
- [59] P. Sułkowski, *Deformed boson-fermion correspondence, Q -bosons, and topological strings on the conifold*, JHEP 0810 (2008) 104 [0808.2327 [hep-th]].
- [60] R. Dijkgraaf, C. Vafa, *Matrix Models, Topological Strings, and Supersymmetric Gauge Theories*, Nucl. Phys. B644 (2002) 3-20 [hep-th/0206255].
- [61] R. Dijkgraaf, C. Vafa, *A Perturbative Window into Non-Perturbative Physics*, hep-th/0208048.
- [62] F. Cachazo, C. Vafa, *$N=1$ and $N=2$ Geometry from Fluxes*, hep-th/0206017.
- [63] T. Hollowood, *Five-Dimensional Gauge Theories and Quantum Mechanical Matrix Models*, JHEP 0303 (2003) 039 [hep-th/0302165].
- [64] M. Wijnholt, *Five-Dimensional Gauge Theories and Unitary Matrix Models*, hep-th/0401025.
- [65] F. Ferrari, *The microscopic approach to $N=1$ super Yang-Mills theories*, Intl. J. Mod. Phys. A23 2307-2323 [0804.0244 [hep-th]].
- [66] So Matsuura, *Instanton Counting and Dielectric Branes*, JHEP 0809 (2008) 083 [0808.3483 [hep-th]].
- [67] Ta-Sheng Tai, *Instanton counting and Matrix Model*, Prog. Theor. Phys. 119 (2008) 165 [0709.0432 [hep-th]].
- [68] P. Byrd, M. Friedman, *Handbook of elliptic integrals for engineers and physicists*, Springer Verlag (1954).